

RANDOM VARIABLES IN A GRAPH W^* -PROBABILITY SPACE

ILWOO CHO

ABSTRACT. In [15], we constructed a W^* -probability space $(W^*(G), E)$ with amalgamation over a von Neumann algebra D_G , where $W^*(G)$ is a graph W^* -algebra induced by the countable directed graph G . In [15], we computed the D_G -valued moments and cumulants of arbitrary random variables in $(W^*(G), E)$ and we could characterize the D_G -freeness of generators of $W^*(G)$, by the so-called diagram-distinctness on G . In this paper, we will observe some special D_G -valued random variables in $(W^*(G), E)$, for instance, D_G -semicircular elements, D_G -even elements, D_G -valued R-diagonal elements and the generating operator of $W^*(G)$. In particular, we can get that (i) if l is a loop in the graph G , then the random variable $L_l + L_l^*$ is D_G -semicircular, (ii) if w is a finite path, then the random variable $L_w + L_w^*$ is D_G -even, (iii) if w is a finite path, then the random variables L_w and L_w^* are D_G -valued R-diagonal.

In this paper, we construct the graph W^* -probability spaces. The graph W^* -probability theory is one of the good example of Speicher's combinatorial free probability theory with amalgamation (See [16]). In this paper, we will observe how to compute the moment and cumulant of an arbitrary random variables in the graph W^* -probability space and the freeness on it with respect to the given conditional expectation. Also, we consider certain special random variables of the graph W^* -probability space, for example, semicircular elements, even elements and R-diagonal elements. This shows that the graph W^* -probability spaces contain the rich free probability objects.

In [10], Kribs and Power defined the free semigroupoid algebras and obtained some properties of them. Our work is highly motivated by [10]. Roughly speaking, graph W^* -algebras are W^* -topology closed version of free semigroupoid algebras. Throughout this paper, let G be a countable directed graph and let $\mathbb{F}^+(G)$ be the free semigroupoid of G , in the sense of Kribs and Power. i.e., it is a collection of all vertices of the graph G as units and all admissible finite paths, under the admissibility. As a set, the free semigroupoid $\mathbb{F}^+(G)$ can be decomposed by

$$\mathbb{F}^+(G) = V(G) \cup FP(G),$$

where $V(G)$ is the vertex set of the graph G and $FP(G)$ is the set of all admissible finite paths. Trivially the edge set $E(G)$ of the graph G is properly contained in

Key words and phrases. Graph W^* -Probability Sapces over the Diagonal Subalgebras, D_G -Freeness, D_G -valued moments and cumulants, D_G -semicircularity, D_G -evenness, D_G -valued R-diagonality, Generating Operators.

$FP(G)$, since all edges of the graph can be regarded as finite paths with their length 1. We define a graph W^* -algebra of G by

$$W^*(G) \stackrel{def}{=} \overline{\mathbb{C}[\{L_w, L_w^* : w \in \mathbb{F}^+(G)\}]}^w,$$

where L_w and L_w^* are creation operators and annihilation operators on the generalized Fock space $H_G = l^2(\mathbb{F}^+(G))$ induced by the given graph G , respectively. Notice that the creation operators induced by vertices are projections and the creation operators induced by finite paths are partial isometries. We can define the W^* -subalgebra D_G of $W^*(G)$, which is called the diagonal subalgebra by

$$D_G \stackrel{def}{=} \overline{\mathbb{C}[\{L_v : v \in V(G)\}]}^w.$$

Then each element a in the graph W^* -algebra $W^*(G)$ is expressed by

$$a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}, \text{ for } p_w \in \mathbb{C},$$

where $\mathbb{F}^+(G : a)$ is a support of the element a , as a subset of the free semigroupoid $\mathbb{F}^+(G)$. The above expression of the random variable a is said to be the Fourier expansion of a . Since $\mathbb{F}^+(G)$ is decomposed by the disjoint subsets $V(G)$ and $FP(G)$, the support $\mathbb{F}^+(G : a)$ of a is also decomposed by the following disjoint subsets,

$$V(G : a) = \mathbb{F}^+(G : a) \cap V(G)$$

and

$$FP(G : a) = \mathbb{F}^+(G : a) \cap FP(G).$$

Thus the operator a can be re-expressed by

$$a = \sum_{v \in V(G:a)} p_v L_v + \sum_{w \in FP(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

Notice that if $V(G : a) \neq \emptyset$, then $\sum_{v \in V(G:a)} p_v L_v$ is contained in the diagonal subalgebra D_G . Thus we have the canonical conditional expectation $E : W^*(G) \rightarrow D_G$, defined by

$$E(a) = \sum_{v \in V(G:a)} p_v L_v,$$

for all $a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}$ in $W^*(G)$. Then the algebraic pair $(W^*(G), E)$ is a W^* -probability space with amalgamation over D_G (See [16]). It is easy to check that the conditional expectation E is faithful in the sense that if $E(a^*a) = 0_{D_G}$, for $a \in W^*(G)$, then $a = 0_{D_G}$.

For the fixed operator $a \in W^*(G)$, the support $\mathbb{F}^+(G : a)$ of the operator a is again decomposed by

$$\mathbb{F}^+(G : a) = V(G : a) \cup FP_*(G : a) \cup FP_*^c(G : a),$$

with the decomposition of $FP(G : a)$,

$$FP(G : a) = FP_*(G : a) \cup FP_*^c(G : a),$$

where

$$FP_*(G : a) = \{w \in FP(G : a) : \text{both } L_w \text{ and } L_w^* \text{ are summands of } a\}$$

and

$$FP_*^c(G : a) = FP(G : a) \setminus FP_*(G : a).$$

The above new expression plays a key role to find the D_G -valued moments of the random variable a . In fact, the summands $p_v L_v$'s and $p_w L_w + p_w^t L_w^*$, for $v \in V(G : a)$ and $w \in FP_*(G : a)$ act for the computation of D_G -valued moments of a . By using the above partition of the support of a random variable, we can compute the D_G -valued moments and D_G -valued cumulants of it via the lattice path model LP_n and the lattice path model LP_n^* satisfying the $*$ -axis-property. At a first glance, the computations of D_G -valued moments and cumulants look so abstract (See Chapter 3) and hence it looks useless. However, these computations, in particular the computation of D_G -valued cumulants, provides us how to figure out the D_G -freeness of random variables by making us compute the mixed cumulants. As applications, in the final chapter, we can compute the moment and cumulant of the operator that is the sum of N -free semicircular elements with their covariance 2. If a is the operator, then the n -th moment of a is

$$\begin{cases} (2N)^{\frac{n}{2}} \cdot c_{\frac{n}{2}} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd,} \end{cases}$$

and the n -th cumulant of a is

$$\begin{cases} 2N & \text{if } n = 2 \\ 0 & \text{otherwise.} \end{cases}$$

Based on the D_G -cumulant computation, we can characterize the D_G -freeness of generators of $W^*(G)$, by the so-called diagram-distinctness on the graph G . i.e., the random variables L_{w_1} and L_{w_2} are free over D_G if and only if w_1 and w_2 are diagram-distinct the sense that w_1 and w_2 have different diagrams on the graph G .

Also, we could find the necessary condition for the D_G -freeness of two arbitrary random variables a and b . i.e., if the supports $\mathbb{F}^+(G : a)$ and $\mathbb{F}^+(G : b)$ are diagram-distinct, in the sense that w_1 and w_2 are diagram distinct for all pairs

$(w_1, w_2) \in \mathbb{F}^+(G : a) \times \mathbb{F}^+(G : b)$, then the random variables a and b are free over D_G .

From Chapter 4 to Chapter 6, we will consider some special D_G -valued random variables in a graph W^* -probability space $(W^*(G), E)$. The those random variables are the basic objects to study Free Probability Theory. We can conclude that

- (i) if l is a loop, then $L_l + L_l^*$ is D_G -semicircular.
- (ii) if w is a finite path, then $L_w + L_w^*$ is D_G -even.
- (iii) if w is a finite path, then L_w and L_w^* are D_G -valued R-diagonal.

In Chapter 5, we consider the generating operator of the graph W^* -algebra $W^*(G)$. We compute the moments and cumulants of the generating operators of the one-vertex graph with N -edges and the circulant graph.

1. GRAPH W^* -PROBABILITY SPACES

Let G be a countable directed graph and let $\mathbb{F}^+(G)$ be the free semigroupoid of G . i.e., the set $\mathbb{F}^+(G)$ is the collection of all vertices as units and all admissible finite paths of G . Let w be a finite path with its source $s(w) = x$ and its range $r(w) = y$, where $x, y \in V(G)$. Then sometimes we will denote w by $w = xwy$ to express the source and the range of w . We can define the graph Hilbert space H_G by the Hilbert space $l^2(\mathbb{F}^+(G))$ generated by the elements in the free semigroupoid $\mathbb{F}^+(G)$. i.e., this Hilbert space has its Hilbert basis $\mathcal{B} = \{\xi_w : w \in \mathbb{F}^+(G)\}$. Suppose that $w = e_1 \dots e_k \in FP(G)$ is a finite path with $e_1, \dots, e_k \in E(G)$. Then we can regard ξ_w as $\xi_{e_1} \otimes \dots \otimes \xi_{e_k}$. So, in [10], Kribs and Power called this graph Hilbert space the generalized Fock space. Throughout this paper, we will call H_G the graph Hilbert space to emphasize that this Hilbert space is induced by the graph.

Define the creation operator L_w , for $w \in \mathbb{F}^+(G)$, by the multiplication operator by ξ_w on H_G . Then the creation operator L on H_G satisfies that

- (i) $L_w = L_{xwy} = L_x L_w L_y$, for $w = xwy$ with $x, y \in V(G)$.

$$(ii) L_{w_1} L_{w_2} = \begin{cases} L_{w_1 w_2} & \text{if } w_1 w_2 \in \mathbb{F}^+(G) \\ 0 & \text{if } w_1 w_2 \notin \mathbb{F}^+(G), \end{cases}$$

for all $w_1, w_2 \in \mathbb{F}^+(G)$.

Now, define the annihilation operator L_w^* , for $w \in \mathbb{F}^+(G)$ by

$$L_w^* \xi_{w'} \stackrel{def}{=} \begin{cases} \xi_h & \text{if } w' = wh \in \mathbb{F}^+(G)\xi \\ 0 & \text{otherwise.} \end{cases}$$

The above definition is gotten by the following observation ;

$$\begin{aligned} \langle L_w \xi_h, \xi_{wh} \rangle &= \langle \xi_{wh}, \xi_{wh} \rangle \\ &= 1 = \langle \xi_h, \xi_h \rangle \\ &= \langle \xi_h, L_w^* \xi_{wh} \rangle, \end{aligned}$$

where $\langle, \rangle$ is the inner product on the graph Hilbert space H_G . Of course, in the above formula we need the admissibility of w and h in $\mathbb{F}^+(G)$. However, even though w and h are not admissible (i.e., $wh \notin \mathbb{F}^+(G)$), by the definition of L_w^* , we have that

$$\begin{aligned} \langle L_w \xi_h, \xi_h \rangle &= \langle 0, \xi_h \rangle \\ &= 0 = \langle \xi_h, 0 \rangle \\ &= \langle \xi_h, L_w^* \xi_h \rangle. \end{aligned}$$

Notice that the creation operator L and the annihilation operator L^* satisfy that

$$(1.1) \quad L_w^* L_w = L_y \quad \text{and} \quad L_w L_w^* = L_x, \quad \text{for all } w = xwy \in \mathbb{F}^+(G),$$

under the weak topology, where $x, y \in V(G)$. Remark that if we consider the von Neumann algebra $W^*(\{L_w\})$ generated by L_w and L_w^* in $B(H_G)$, then the projections L_y and L_x are Murray-von Neumann equivalent, because there exists a partial isometry L_w satisfying the relation (1.1). Indeed, if $w = xwy$ in $\mathbb{F}^+(G)$, with $x, y \in V(G)$, then under the weak topology we have that

$$(1.2) \quad L_w L_w^* L_w = L_w \quad \text{and} \quad L_w^* L_w L_w^* = L_w^*.$$

So, the creation operator L_w is a partial isometry in $W^*(\{L_w\})$ in $B(H_G)$. Assume now that $v \in V(G)$. Then we can regard v as $v = vvv$. So,

$$(1.3) \quad L_v^* L_v = L_v = L_v L_v^* = L_v^*.$$

This relation shows that L_v is a projection in $B(H_G)$ for all $v \in V(G)$.

Define the **graph W^* -algebra** $W^*(G)$ by

$$W^*(G) \stackrel{def}{=} \overline{\mathbb{C}[\{L_w, L_w^* : w \in \mathbb{F}^+(G)\}]}^w.$$

Then all generators are either partial isometries or projections, by (1.2) and (1.3). So, this graph W^* -algebra contains a rich structure, as a von Neumann algebra. (This construction can be the generalization of that of group von Neumann algebra.)

Naturally, we can define a von Neumann subalgebra $D_G \subset W^*(G)$ generated by all projections L_v , $v \in V(G)$. i.e.

$$D_G \stackrel{def}{=} W^*(\{L_v : v \in V(G)\}).$$

We call this subalgebra the **diagonal subalgebra** of $W^*(G)$. Notice that $D_G = \Delta_{|G|} \subset M_{|G|}(\mathbb{C})$, where $\Delta_{|G|}$ is the subalgebra of $M_{|G|}(\mathbb{C})$ generated by all diagonal matrices. Also, notice that $1_{D_G} = \sum_{v \in V(G)} L_v = 1_{W^*(G)}$.

If $a \in W^*(G)$ is an operator, then it has the following decomposition which is called the Fourier expansion of a ;

$$(1.4) \quad a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1, *\}} p_w L_w^{u_w},$$

where $p_w \in \mathbb{C}$ and $\mathbb{F}^+(G:a)$ is the support of a defined by

$$\mathbb{F}^+(G:a) = \{w \in \mathbb{F}^+(G) : p_w \neq 0\}.$$

Remark that the free semigroupoid $\mathbb{F}^+(G)$ has its partition $\{V(G), FP(G)\}$, as a set. i.e.,

$$\mathbb{F}^+(G) = V(G) \cup FP(G) \quad \text{and} \quad V(G) \cap FP(G) = \emptyset.$$

So, the support of a is also partitioned by

$$\mathbb{F}^+(G:a) = V(G:a) \cup FP(G:a),$$

where

$$V(G:a) \stackrel{def}{=} V(G) \cap \mathbb{F}^+(G:a)$$

and

$$FP(G:a) \stackrel{def}{=} FP(G) \cap \mathbb{F}^+(G:a).$$

So, the above Fourier expansion (1.4) of the random variable a can be re-expressed by

$$(1.5) \quad a = \sum_{v \in V(G:a)} p_v L_v + \sum_{w \in FP(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

We can easily see that if $V(G:a) \neq \emptyset$, then $\sum_{v \in V(G:a)} p_v L_v$ is contained in the diagonal subalgebra D_G . Also, if $V(G:a) = \emptyset$, then $\sum_{v \in V(G:a)} p_v L_v = 0_{D_G}$. So, we can define the following canonical conditional expectation $E : W^*(G) \rightarrow D_G$ by

$$(1.6) \quad E(a) = E \left(\sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1,*\}} p_w L_w^{u_w} \right) \stackrel{def}{=} \sum_{v \in V(G:a)} p_v L_v,$$

for all $a \in W^*(G)$. Indeed, E is a well-determined conditional expectation ; it is a bimodule map satisfying that

$$E(d) = d, \text{ for all } d \in D_G.$$

And

$$\begin{aligned} E(dad') &= E(d(a_d + a_0)d') = E(da_d d' + da_0 d') \\ &= E(da_d d') = da_d d' = d(E(a))d', \end{aligned}$$

for all $d, d' \in D_G$ and $a = a_d + a_0 \in W^*(G)$, where

$$a_d = \sum_{v \in V(G:a)} p_v L_v \quad \text{and} \quad a_0 = \sum_{w \in FP(G:a), u_w \in \{1,*\}} p_w L_w^{u_w}.$$

Also,

$$E(a^*) = E((a_d + a_0)^*) = E(a_d^* + a_0^*) = a_d^* = E(a)^*,$$

for all $a \in W^*(G)$. Here, $a_d^* = \left(\sum_{v \in V(G:a)} p_v L_v \right)^* = \sum_{v \in V(G:a)} \overline{p_v} L_v$ in D_G .

Definition 1.1. Let G be a countable directed graph and let $W^*(G)$ be the graph W^* -algebra induced by G . Let $E : W^*(G) \rightarrow D_G$ be the conditional expectation defined above. Then we say that the algebraic pair $(W^*(G), E)$ is the graph W^* -probability space over the diagonal subalgebra D_G . By the very definition, it is one of the W^* -probability space with amalgamation over D_G . All elements in $(W^*(G), E)$ are called D_G -valued random variables.

We have a graph W^* -probability space $(W^*(G), E)$ over its diagonal subalgebra D_G . We will define the following free probability data of D_G -valued random variables.

Definition 1.2. Let $W^*(G)$ be the graph W^* -algebra induced by G and let $a \in W^*(G)$. Define the n -th (D_G -valued) moment of a by

$$E(d_1 a d_2 a \dots d_n a), \text{ for all } n \in \mathbb{N},$$

where $d_1, \dots, d_n \in D_G$. Also, define the n -th (D_G -valued) cumulant of a by

$$k_n(d_1 a, d_2 a, \dots, d_n a) = C^{(n)}(d_1 a \otimes d_2 a \otimes \dots \otimes d_n a),$$

for all $n \in \mathbb{N}$, and for $d_1, \dots, d_n \in D_G$, where $\widehat{C} = (C^{(n)})_{n=1}^\infty \in I^c(W^*(G), D_G)$ is the cumulant multiplicative bimodule map induced by the conditional expectation E , in the sense of Speicher. We define the n -th trivial moment of a and the n -th trivial cumulant of a by

$$E(a^n) \quad \text{and} \quad k_n \left(\underbrace{a, a, \dots, a}_{n\text{-times}} \right) = C^{(n)}(a \otimes a \otimes \dots \otimes a),$$

respectively, for all $n \in \mathbb{N}$.

To compute the D_G -valued moments and cumulants of the D_G -valued random variable a , we need to introduce the following new definition ;

Definition 1.3. Let $(W^*(G), E)$ be a graph W^* -probability space over D_G and let $a \in (W^*(G), E)$ be a random variable. Define the subset $FP_*(G : a)$ in $FP(G : a)$ by

$$FP_*(G : a) \stackrel{\text{def}}{=} \{w \in \mathbb{F}^+(G : a) : \text{both } L_w \text{ and } L_w^* \text{ are summands of } a\}.$$

And let $FP_*^c(G : a) \stackrel{\text{def}}{=} FP(G : a) \setminus FP_*(G : a)$.

We already observed that if $a \in (W^*(G), E)$ is a D_G -valued random variable, then a has its Fourier expansion $a_d + a_0$, where

$$a_d = \sum_{v \in V(G:a)} p_v L_v$$

and

$$a_0 = \sum_{w \in FP(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

By the previous definition, the set $FP(G : a)$ is partitioned by

$$FP(G : a) = FP_*(G : a) \cup FP_*^c(G : a),$$

for the fixed random variable a in $(W^*(G), E)$. So, the summand a_0 , in the Fourier expansion of $a = a_d + a_0$, has the following decomposition ;

$$a_0 = a_{(*)} + a_{(non-*)},$$

where

$$a_{(*)} = \sum_{l \in FP_*(G:a)} (p_l L_l + p_{l^t} L_l^*)$$

and

$$a_{(non-*)} = \sum_{w \in FP_*(G:a), u_w \in \{1,*\}} p_w L_w^{u_w},$$

where p_{l^t} is the coefficient of L_l^* depending on $l \in FP_*(G : a)$. (There is no special meaning for the complex number p_{l^t} . But we have to keep in mind that $p_l \neq p_{l^t}$, in general. i.e. $a_{(*)} = \sum_{l_1 \in FP_*(G:a)} p_{l_1} L_{l_1} + \sum_{l_2 \in FP_*(G:a)} p_{l_2} L_{l_2}^*$! But for the convenience of using notation, we will use the notation p_{l^t} , for the coefficient of L_l^* .) For instance, let $V(G : a) = \{v_1, v_2\}$ and $FP(G : a) = \{w_1, w_2\}$ and let the random variable a in $(W^*(G), E)$ be

$$a = L_{v_1} + L_{v_2} + L_{w_1}^* + L_{w_1} + L_{w_2}^*.$$

Then we have that $a_d = L_{v_1} + L_{v_2}$, $a_{(*)} = L_{w_1}^* + L_{w_1}$ and $a_{(non-*)} = L_{w_2}^*$. By definition, $a_0 = a_{(*)} + a_{(non-*)}$.

2. D_G -MOMENTS AND D_G -CUMULANTS OF RANDOM VARIABLES

Throughout this chapter, let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over its diagonal subalgebra D_G . In this chapter, we will compute the D_G -valued moments and the D_G -valued cumulants of arbitrary random variable

$$a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1,*\}} p_w L_w^{u_w}$$

in the graph W^* -probability space $(W^*(G), E)$.

2.1. Lattice Path Model.

Throughout this section, let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over its diagonal subalgebra D_G . Let $w_1, \dots, w_n \in \mathbb{F}^+(G)$ and let $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \in (W^*(G), E)$ be a D_G -valued random variable. In this section, we will define a lattice path model for the random variable $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$. Recall that if $w = e_1 \dots e_k \in FP(G)$ with $e_1, \dots, e_k \in E(G)$, then we can define the length $|w|$ of w by k . i.e.e, the length $|w|$ of w is the cardinality k of the admissible edges $e_1, \dots, e_k$.

Definition 2.1. Let G be a countable directed graph and $\mathbb{F}^+(G)$, the free semi-groupoid. If $w \in \mathbb{F}^+(G)$, then L_w is the corresponding D_G -valued random variable in $(W^*(G), E)$. We define the lattice path l_w of L_w and the lattice path l_w^{-1} of L_w^* by the lattice paths satisfying that ;

- (i) the lattice path l_w starts from $*$ $= (0, 0)$ on the $\mathbb{R}^2$ -plane.
- (ii) if $w \in V(G)$, then l_w has its end point $(0, 1)$.
- (iii) if $w \in E(G)$, then l_w has its end point $(1, 1)$.
- (iv) if $w \in E(G)$, then l_w^{-1} has its end point $(-1, -1)$.
- (v) if $w \in FP(G)$ with $|w| = k$, then l_w has its end point (k, k) .
- (vi) if $w \in FP(G)$ with $|w| = k$, then l_w^{-1} has its end point $(-k, -k)$.

Assume that finite paths $w_1, \dots, w_s$ in $FP(G)$ satisfy that $w_1 \dots w_s \in FP(G)$. Define the lattice path $l_{w_1 \dots w_s}$ by the connected lattice path of the lattice paths $l_{w_1}, \dots, l_{w_s}$. i.e.e, l_{w_2} starts from $(k_{w_1}, k_{w_1}) \in \mathbb{R}^+$ and ends at $(k_{w_1} + k_{w_2}, k_{w_1} + k_{w_2})$, where $|w_1| = k_{w_1}$ and $|w_2| = k_{w_2}$. Similarly, we can define the lattice path $l_{w_1 \dots w_s}^{-1}$ as the connected path of $l_{w_s}^{-1}, l_{w_{s-1}}^{-1}, \dots, l_{w_1}^{-1}$.

Definition 2.2. Let G be a countable directed graph and assume that $L_{w_1}, \dots, L_{w_n}$ are generators of $(W^*(G), E)$. Then we have the lattice paths $l_{w_1}, \dots, l_{w_n}$ of $L_{w_1}, \dots, L_{w_n}$, respectively in $\mathbb{R}^2$. Suppose that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \neq 0_{D_G}$ in $(W^*(G), E)$, where $u_{w_1}, \dots, u_{w_n} \in \{1, *\}$. Define the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ of nonzero $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ by the connected lattice path of $l_{w_1}^{t_{w_1}}, \dots, l_{w_n}^{t_{w_n}}$, where $t_{w_j} = 1$ if $u_{w_j} = 1$ and $t_{w_j} = -1$ if $u_{w_j} = *$. Assume that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = 0_{D_G}$. Then the empty set $\emptyset$ in $\mathbb{R}^2$ is the lattice path of it. We call it the empty lattice path. By LP_n , we will denote the set of all lattice paths of the D_G -valued random variables having their forms of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$, including empty lattice path.

Also, we will define the following important property on the set of all lattice paths ;

Definition 2.3. Let $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \neq \emptyset$ be a lattice path of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \neq 0_{D_G}$ in LP_n . If the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ starts from $*$ and ends on the $*$ -axis in $\mathbb{R}^+$, then we say that the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ has the $*$ -axis-property. By LP_n^* , we will denote the set of all lattice paths having their forms of $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ which have the $*$ -axis-property. By little abuse of notation, sometimes, we will say that the D_G -valued random variable $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ satisfies the $*$ -axis-property if the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ of it has the $*$ -axis-property.

The following theorem shows that finding $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}})$ is checking the $*$ -axis-property of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$.

Theorem 2.1. *Let $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \in (W^*(G), E)$ be a D_G -valued random variable, where $u_{w_1}, \dots, u_{w_n} \in \{1, *\}$. Then $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}) \neq 0_{D_G}$ if and only if $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ has the $*$ -axis-property (i.e., the corresponding lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ is contained in LP_n^* . Notice that $\emptyset \notin LP_n^*$.)*

Proof. ($\Leftarrow$) Let $l = l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n^*$. Suppose that $w_1 = vw_1v'_1$ and $w_n = v_nw_nv'_n$, for $v_1, v'_1, v_n, v'_n \in V(G)$. If l is in LP_n^* , then

$$(2.1.1) \quad \begin{cases} v_1 = v'_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = 1 \\ v_1 = v_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = * \\ v'_1 = v'_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = 1 \\ v'_1 = v_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = *. \end{cases}$$

By the definition of lattice paths having the $*$ -axis-property and by (2.1.1), if $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n^*$, then there exists $v \in V(G)$ such that

$$(2.1.2) \quad \begin{cases} L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = L_v, \\ \text{where} \\ \begin{cases} v = v_1 = v'_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = 1 \\ v = v_1 = v_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = * \\ v = v'_1 = v'_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = 1 \\ v = v'_1 = v_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = *. \end{cases} \end{cases}$$

This shows that $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}) = L_v \neq 0_{D_G}$.

($\Rightarrow$) Assume that $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}) \neq 0_{D_G}$. This means that there exists L_v , with $v \in V(G)$, such that

$$(2.1.3) \quad E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}) = L_v.$$

Equivalently, we have that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = L_v$ in $W^*(G)$. Let $l = l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n$ be the lattice path of the D_G -valued random variable $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$. By (2.1.3), trivially, $l \neq \emptyset$, since l should be the connected lattice path. Assume that the nonempty lattice path l is contained in $LP_n \setminus LP_n^*$. Then, under the same conditions of (2.1.1), we have that

$$(2.1.4) \quad \begin{cases} v_1 \neq v'_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = 1 \\ v_1 \neq v_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = * \\ v'_1 \neq v'_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = 1 \\ v'_1 \neq v_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = *. \end{cases}$$

Therefore, by (2.1.2), there is no vertex v satisfying $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = L_v$. This contradicts our assumption. ■

By the previous theorem, we can conclude that $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}) = L_v$, for some $v \in V(G)$ if and only if the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ has the $*$ -axis-property (i.e., $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n^*$).

2.2. D_G -Valued Moments and Cumulants of Random Variables. Let $w_1, \dots, w_n \in \mathbb{F}^+(G)$, $u_1, \dots, u_n \in \{1, *\}$ and let $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \in (W^*(G), E)$ be a D_G -valued random variable. Recall that, in the previous section, we observed that the D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} = L_v \in (W^*(G), E)$ with $v \in V(G)$ if and only if the lattice path $l_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ of $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ has the $*$ -axis-property (equivalently, $l_{w_1, \dots, w_n}^{u_1, \dots, u_n} \in LP_n^*$). Throughout this section, fix a D_G -valued random variable $a \in (W^*(G), E)$. Then the D_G -valued random variable a has the following Fourier expansion,

$$a = \sum_{v \in V(G:a)} p_v L_v + \sum_{l \in FP_*(G:a)} (p_l L_l + p_{l^t} L_{l^t}) + \sum_{w \in FPC_*(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

Let's observe the new D_G -valued random variable $d_1 a d_2 a \dots d_n a \in (W^*(G), E)$, where $d_1, \dots, d_n \in D_G$ and $a \in W^*(G)$ is given. Put

$$d_j = \sum_{v_j \in V(G:d_j)} q_{v_j} L_{v_j} \in D_G, \text{ for } j = 1, \dots, n.$$

Notice that $V(G:d_j) = \mathbb{F}^+(G:d_j)$, since $d_j \in D_G \hookrightarrow W^*(G)$. Then

$$\begin{aligned} & d_1 a d_2 a \dots d_n a \\ &= \left(\sum_{v_1 \in V(G:d_1)} q_{v_1} L_{v_1} \right) \left(\sum_{w_1 \in \mathbb{F}^+(G:a), u_{w_1} \in \{1, *\}} p_{w_1} L_{w_1}^{u_{w_1}} \right) \\ & \quad \dots \left(\sum_{v_n \in V(G:d_n)} q_{v_n} L_{v_n} \right) \left(\sum_{w_n \in \mathbb{F}^+(G:a), u_{w_n} \in \{1, *\}} p_{w_n} L_{w_n}^{u_{w_n}} \right) \\ &= \sum_{(v_1, \dots, v_n) \in \prod_{j=1}^n V(G:d_j)} (q_{v_1} \dots q_{v_n}) \end{aligned}$$

$$(1.2.1) \quad \left(L_{v_1} \left(\sum_{w_1 \in \mathbb{F}^+(G:a), u_{w_1} \in \{1, *\}} p_{w_1} L_{w_1}^{u_{w_1}} \right) \right. \\ \left. \cdots L_{v_n} \left(\sum_{w_n \in \mathbb{F}^+(G:a), u_{w_n} \in \{1, *\}} p_{w_n} L_{w_n}^{u_{w_n}} \right) \right)$$

$$= \sum_{(v_1, \dots, v_n) \in \Pi_{j=1}^n V(G:d_j)} (q_{v_1} \cdots q_{v_n}) \\ \sum_{(w_1, \dots, w_n) \in \mathbb{F}^+(G:a)^n, u_{w_j} \in \{1, *\}} (p_{w_1} \cdots p_{w_n}) L_{v_1}^{u_{w_1}} \cdots L_{v_n}^{u_{w_n}}.$$

Now, consider the random variable $L_{v_1}^{u_{w_1}} \cdots L_{v_n}^{u_{w_n}}$ in the formula (1.2.1). Suppose that $w_j = x_j w_j y_j$, with $x_j, y_j \in V(G)$, for all $j = 1, \dots, n$. Then

$$(1.2.2) \quad L_{v_1}^{u_{w_1}} \cdots L_{v_n}^{u_{w_n}} = \delta_{(v_1, x_1, y_1: u_{w_1})} \cdots \delta_{(v_n, x_n, y_n: u_{w_n})} (L_{w_1}^{u_{w_1}} \cdots L_{w_n}^{u_{w_n}}),$$

where

$$\delta_{(v_j, x_j, y_j: u_{w_j})} = \begin{cases} \delta_{v_j, x_j} & \text{if } u_{w_j} = 1 \\ \delta_{v_j, y_j} & \text{if } u_{w_j} = *, \end{cases}$$

for all $j = 1, \dots, n$, where δ in the right-hand side is the Kronecker delta. So, the left-hand side can be understood as a (conditional) Kronecker delta depending on $\{1, *\}$.

By (1.2.1) and (1.2.2), the n -th moment of a is

$$E(d_1 a \cdots d_n a) \\ = E \left(\sum_{(v_1, \dots, v_n) \in \Pi_{j=1}^n V(G:d_j)} (\Pi_{j=1}^n q_{v_j}) \right. \\ \sum_{(w_1, \dots, w_n) \in \mathbb{F}^+(G:a)^n, w_j = x_j w_j y_j, u_{w_j} \in \{1, *\}} (\Pi_{j=1}^n p_{w_j}) \\ \left. \left(\Pi_{j=1}^n \delta_{(v_j, x_j, y_j: u_{w_j})} \right) (L_{w_1}^{u_{w_1}} \cdots L_{w_n}^{u_{w_n}}) \right) \\ = \sum_{(v_1, \dots, v_n) \in \Pi_{j=1}^n V(G:d_j)} (\Pi_{j=1}^n q_{v_j})$$

$$\sum_{(w_1, \dots, w_n) \in \mathbb{F}^+(G:a)^n, w_j = x_j w_j y_j, u_{w_j} \in \{1, *\}} \left(\prod_{j=1}^n p_{w_j} \right) \\ \left(\prod_{j=1}^n \delta_{(v_j, x_j, y_j : u_{w_j})} \right) E \left(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \right).$$

Thus to compute the n -th moment of a , we have to observe $E \left(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \right)$. In the previous section, we observed that $E \left(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \right)$ is nonvanishing if and only if $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ has the $*$ -axis-property.

Proposition 2.2. *Let $a \in (W^*(G), E)$ be given as above. Then the n -th moment of a is*

$$E(d_1 a \dots d_n a) = \sum_{(v_1, \dots, v_n) \in \prod_{j=1}^n V(G:d_j)} \left(\prod_{j=1}^n q_{v_j} \right) \\ \sum_{(w_1, \dots, w_n) \in \mathbb{F}^+(G:a)^n, u_{w_j} \in \{1, *\}, l_{w_1}^{u_{w_1}}, \dots, l_{w_n}^{u_{w_n}} \in LP_n^*} \left(\prod_{j=1}^n p_{w_j} \right) \\ \left(\prod_{j=1}^n \delta_{(v_j, x_j, y_j : u_{w_j})} \right) E \left(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \right).$$

□

From now, rest of this paper, we will compute the D_G -valued cumulants of the given D_G -valued random variable a . Let $w_1, \dots, w_n \in FP(G)$ be finite paths and $u_1, \dots, u_n \in \{1, *\}$. Then, by the Möbius inversion, we have

$$(2.2.1) \quad k_n \left(L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n} \right) = \sum_{\pi \in NC(n)} \widehat{E}(\pi) \left(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n} \right) \mu(\pi, 1_n),$$

where $\widehat{E} = (E^{(n)})_{n=1}^\infty$ is the moment multiplicative bimodule map induced by the conditional expectation E (See [16]) and where $NC(n)$ is the collection of all noncrossing partition over $\{1, \dots, n\}$. Notice that if $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ does not have the $*$ -axis-property, then

$$E \left(L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \right) = 0_{D_G},$$

by Section 2.1. Consider the noncrossing partition $\pi \in NC(n)$ with its blocks $V_1, \dots, V_k$. Choose one block $V_j = (j_1, \dots, j_k) \in \pi$. Then we have that

$$(2.2.2) \quad \widehat{E}(\pi |_{V_j}) \left(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n} \right) = E \left(L_{w_{j_1}}^{u_{j_1}} d_{j_2} L_{w_{j_2}}^{u_{j_2}} \dots d_{j_k} L_{w_{j_k}}^{u_{j_k}} \right),$$

where

$$d_{j_i} = \begin{cases} 1_{D_G} & \text{if there is no inner blocks} \\ & \text{between } j_{i-1} \text{ and } j_i \text{ in } V_j \\ L_{v_{j_i}} \neq 1_{D_G} & \text{if there are inner blocks} \\ & \text{between } j_{i-1} \text{ and } j_i \text{ in } V_j, \end{cases}$$

where $v_{j_2}, \dots, v_{j_k} \in V(G)$. So, again by Section 2.1, $\widehat{E}(\pi |_{V_j})(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n})$ is nonvanishing if and only if $L_{w_{j_1}}^{u_{j_1}} d_{j_2} L_{w_{j_2}}^{u_{j_2}} \dots d_{j_k} L_{w_{j_k}}^{u_{j_k}}$ has the $*$ -axis-property, for all $j = 1, \dots, n$.

Assume that

$$\widehat{E}(\pi |_{V_j})(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) = L_{v_j}$$

and

$$\widehat{E}(\pi |_{V_i})(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) = L_{v_i}.$$

If $v_j \neq v_i$, then the partition-dependent D_G -moment satisfies that

$$\widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) = 0_{D_G}.$$

This says that $\widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) \neq 0_{D_G}$ if and only if there exists $v \in V(G)$ such that

$$\widehat{E}(\pi |_{V_j})(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) = L_v,$$

for all $j = 1, \dots, k$.

Definition 2.4. Let $NC(n)$ be the set of all noncrossing partition over $\{1, \dots, n\}$ and let $L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n} \in (W^*(G), E)$ be D_G -valued random variables, where $u_1, \dots, u_n \in \{1, *\}$. We say that the D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ is π -connected if the π -dependent D_G -moment of it is nonvanishing, for $\pi \in NC(n)$. In other words, the random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ is π -connected, for $\pi \in NC(n)$, if

$$\widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) \neq 0_{D_G}.$$

i.e., there exists a vertex $v \in V(G)$ such that

$$\widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) = L_v.$$

For convenience, we will define the following subset of $NC(n)$;

Definition 2.5. Let $NC(n)$ be the set of all noncrossing partitions over $\{1, \dots, n\}$ and fix a D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ in $(W^*(G), E)$, where $u_1, \dots, u_n \in \{1, *\}$. For the fixed D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$, define

$$C_{w_1, \dots, w_n}^{u_1, \dots, u_n} \stackrel{\text{def}}{=} \{\pi \in NC(n) : L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \text{ is } \pi\text{-connected}\},$$

in $NC(n)$. Let μ be the Möbius function in the incidence algebra I_2 . Define the number $\mu_{w_1, \dots, w_n}^{u_1, \dots, u_n}$, for the fixed D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$, by

$$\mu_{w_1, \dots, w_n}^{u_1, \dots, u_n} \stackrel{\text{def}}{=} \sum_{\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}} \mu(\pi, 1_n).$$

Assume that there exists $\pi \in NC(n)$ such that $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} = L_v$ is π -connected. Then $\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ and there exists the maximal partition $\pi_0 \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ such that $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} = L_v$ is π_0 -connected. (Recall that $NC(n)$ is a lattice. We can restrict this lattice ordering on $C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ and hence it is a POset, again.) Notice that $1_n \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$. Therefore, the maximal partition in $C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ is 1_n . Hence we have that ;

Lemma 2.3. *Let $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \in (W^*(G), E)$ be a D_G -valued random variable having the $*$ -axis-property. Then*

$$E(L_{w_1}^{u_1} \dots L_{w_n}^{u_n}) = \widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}),$$

for all $\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$.

Proof. By the previous discussion, we can get the result. ■

By the previous lemmas, we have that

Theorem 2.4. *Let $n \in 2\mathbb{N}$ and let $L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n} \in (W^*(G), E)$ be D_G -valued random variables, where $w_1, \dots, w_n \in FP(G)$ and $u_j \in \{1, *\}$, $j = 1, \dots, n$. Then*

$$k_n(L_{w_1}^{u_1} \dots L_{w_n}^{u_n}) = \mu_{w_1, \dots, w_n}^{u_1, \dots, u_n} \cdot E(L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n}),$$

$$\text{where } \mu_{w_1, \dots, w_n}^{u_1, \dots, u_n} = \sum_{\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}} \mu(\pi, 1_n).$$

Proof. We can compute that

$$\begin{aligned} k_n(L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n}) &= \sum_{\pi \in NC(n)} \widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) \mu(\pi, 1_n) \\ &= \sum_{\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}} \widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) \mu(\pi, 1_n) \end{aligned}$$

by the π -connectedness

$$= \sum_{\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}} E(L_{w_1}^{u_1} \dots L_{w_n}^{u_n}) \mu(\pi, 1_n)$$

by the previous lemma

$$= \left(\sum_{\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}} \mu(\pi, 1_n) \right) \cdot E(L_{w_1}^{u_1} \dots L_{w_n}^{u_n}).$$

■

Now, we can get the following D_G -valued cumulants of the random variable a ;

Corollary 2.5. *Let $n \in \mathbb{N}$ and let $a = a_d + a_{(*)} + a_{(non-*)} \in (W^*(G), E)$ be our D_G -valued random variable. Then $k_1(d_1 a) = d_1 a_d$ and $k_n(d_1 a, \dots, d_n a) = 0_{D_G}$, for all odd n . If $n \in \mathbb{N} \setminus \{1\}$, then*

$$\begin{aligned} k_n(d_1 a, \dots, d_n a) &= \sum_{(v_1, \dots, v_n) \in \prod_{j=1}^n V(G; d_j)} \left(\prod_{j=1}^n q_{v_j} \right) \\ &\quad \sum_{(w_1, \dots, w_n) \in FP_*(G; a)^n, w_j = x_j w_j y_j, u_{w_j} \in \{1, *\}, l_{w_1, \dots, w_n}^{u_1, \dots, u_n} \in LP_n^*} \left(\prod_{j=1}^n p_{w_j} \right) \\ &\quad \left(\prod_{j=1}^n \delta_{(v_j, x_j, y_j; u_j)} \right) \left(\mu_{w_1, \dots, w_n}^{u_1, \dots, u_n} \cdot E(L_{w_1}^{u_1} \dots L_{w_n}^{u_n}) \right), \end{aligned}$$

where $d_1, \dots, d_n \in D_G$ are arbitrary. $\square$

We have the following trivial D_G -valued moments and cumulants of an arbitrary D_G -valued random variable ;

Corollary 2.6. *Let $a \in (W^*(G), E)$ be a D_G -valued random variable and let $n \in \mathbb{N}$. Then*

(1) *The n -th trivial D_G -valued moment of a is*

$$\begin{aligned} E(a^n) &= \sum_{(w_1, \dots, w_n) \in \mathbb{F}^+(G; a)^n, u_{w_j} \in \{1, *\}, l_{w_1, \dots, w_n}^{u_1, \dots, u_n} \in LP_n^*} \\ &\quad \left(\prod_{j=1}^n p_{w_j} \right) \cdot E(L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n}). \end{aligned}$$

(2) *The n -th trivial D_G -valued cumulant of a is*

$$k_1(a) = E(a) = a_d$$

and

$$k_n \left(\underbrace{a, \dots, a}_{n\text{-times}} \right) = \sum_{(w_1, \dots, w_n) \in FP_*(G:a)^n, u_{w_j} \in \{1, *\}, l_{w_1}^{u_1}, \dots, l_{w_n}^{u_n} \in LP_n^*} \left(\prod_{j=1}^n p_{w_j} \right) \left(\mu_{w_1, \dots, w_n}^{u_1, \dots, u_n} \cdot E \left(L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n} \right) \right),$$

where $d_1, \dots, d_n \in D_G$ are arbitrary. $\square$

3. D_G -FREEDOM ON $(W^*(G), E)$

Like before, throughout this chapter, let G be a countable directed graph and $(W^*(G), E)$, the graph W^* -probability space over its diagonal subalgebra D_G . In this chapter, we will consider the D_G -valued freeness of given two random variables in $(W^*(G), E)$. We will characterize the D_G -freeness of D_G -valued random variables L_{w_1} and L_{w_2} , where $w_1 \neq w_2 \in FP(G)$. And then we will observe the D_G -freeness of arbitrary two D_G -valued random variables a_1 and a_2 in terms of their supports. Let

$$(3.0) \quad a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1, *\}} p_w L_w^{u_w} \quad \& \quad b = \sum_{w' \in \mathbb{F}^+(G:b), u_{w'} \in \{1, *\}} p_{w'} L_{w'}^{u_{w'}}$$

be fixed D_G -valued random variables in $(W^*(G), E)$.

Now, fix $n \in \mathbb{N}$ and let $(a_{i_1}^{\varepsilon_{i_1}}, \dots, a_{i_n}^{\varepsilon_{i_n}}) \in \{a, b, a^*, b^*\}^n$, where $\varepsilon_{i_j} \in \{1, *\}$. For convenience, put

$$a_{i_j}^{\varepsilon_{i_j}} = \sum_{w_{i_j} \in \mathbb{F}^+(G:a), u_{i_j} \in \{1, *\}} p_{w_{i_j}}^{(j)} L_{w_{i_j}}^{u_{i_j}}, \text{ for } j = 1, \dots, n.$$

Then, by the little modification of Section , we have that ;

(3.1)

$$\begin{aligned} & E \left(d_{i_1} a_{i_1}^{\varepsilon_{i_1}} \dots d_{i_n} a_{i_n}^{\varepsilon_{i_n}} \right) \\ &= \sum_{(v_{i_1}, \dots, v_{i_n}) \in \prod_{k=1}^n V(G:d_{i_k})} \left(\prod_{k=1}^n q_{v_{i_k}} \right) \end{aligned}$$

$$\sum_{(w_{i_1}, \dots, w_{i_n}) \in \Pi_{k=1}^n \mathbb{F}^+(G: a_{i_k}), w_{i_j} = x_{i_j} w_{i_j} y_{i_j}, u_{i_j} \in \{1, *\}} \left(\prod_{k=1}^n p_{w_{i_k}}^{(k)} \right) \\ \left(\prod_{j=1}^n \delta_{(v_{i_j}, x_{i_j}, y_{i_j}: u_{i_j})} \right) E \left(L_{w_{i_1}}^{u_{i_1}} \dots L_{w_{i_n}}^{u_{i_n}} \right).$$

Therefore, we have that

(3.2)

$$k_n \left(d_{i_1} a_{i_1}^{\varepsilon_{i_1}}, \dots, d_{i_n} a_{i_n}^{\varepsilon_{i_n}} \right) = \sum_{(v_{i_1}, \dots, v_{i_n}) \in \Pi_{k=1}^n V(G: d_{i_k})} \left(\prod_{k=1}^n q_{v_{i_k}} \right) \\ \sum_{(w_{i_1}, \dots, w_{i_n}) \in \Pi_{k=1}^n FP(G: a_{i_k}), w_{i_j} = x_{i_j} w_{i_j} y_{i_j}, u_{i_j} \in \{1, *\}} \left(\prod_{k=1}^n p_{w_{i_k}}^{(k)} \right) \\ \left(\prod_{j=1}^n \delta_{(v_{i_j}, x_{i_j}, y_{i_j}: u_{i_j})} \right) \left(\mu_{w_{i_1}, \dots, w_{i_n}}^{u_{i_1}, \dots, u_{i_n}} \cdot E \left(L_{w_{i_1}}^{u_{i_1}} \dots L_{w_{i_n}}^{u_{i_n}} \right) \right),$$

where $\mu_{w_{i_1}, \dots, w_{i_n}}^{u_{i_1}, \dots, u_{i_n}} = \sum_{\pi \in C_{w_{i_1}, \dots, w_{i_n}}^{u_{i_1}, \dots, u_{i_n}}} \mu(\pi, 1_n)$ and

$$C_{w_{i_1}, \dots, w_{i_n}}^{u_{i_1}, \dots, u_{i_n}} = \{ \pi \in NC^{(even)}(n) : L_{w_{i_1}}^{u_{i_1}} \dots L_{w_{i_n}}^{u_{i_n}} \text{ is } \pi\text{-connected} \}.$$

So, we have the following proposition, by the straightforward computation ;

Proposition 3.1. *Let $a, b \in (W^*(G), E)$ be D_G -valued random variables, such that $a \notin W^*(\{b\}, D_G)$, and let $(a_{i_1}^{\varepsilon_{i_1}}, \dots, a_{i_n}^{\varepsilon_{i_n}}) \in \{a, b, a^*, b^*\}^n$, for $n \in \mathbb{N} \setminus \{1\}$, where $\varepsilon_{i_j} \in \{1, *\}$, $j = 1, \dots, n$. Then*

(3.3)

$$k_n \left(d_{i_1} a_{i_1}^{\varepsilon_{i_1}}, \dots, d_{i_n} a_{i_n}^{\varepsilon_{i_n}} \right) \\ = \sum_{(v_1, \dots, v_n) = (x, y, \dots, x, y) \in \Pi_{j=1}^n V(G: d_j)} \left(\prod_{j=1}^n q_{v_j} \right) \\ \sum_{(w_{i_1}, \dots, w_{i_n}) \in \left(\prod_{k=1}^n FP_*(G: a_{i_k}) \right) \cup W_*^{i w_1, \dots, w_n}, w_{i_j} = x_{i_j} w_{i_j} y_{i_j}} \left(\prod_{k=1}^n p_{w_{i_k}}^{(k)} \right) \\ \left(\prod_{j=1}^n \delta_{(v_{i_j}, x_{i_j}, y_{i_j}: u_{i_j})} \right) \left(\mu_{w_{i_1}, \dots, w_{i_n}}^{u_{i_1}, \dots, u_{i_n}} \cdot \text{Pr oj} \left(L_{w_{i_1}}^{u_{i_1}} \dots L_{w_{i_n}}^{u_{i_n}} \right) \right)$$

where $\mu_n = \sum_{\pi \in C_{w_{i_1}, \dots, w_{i_n}}^{u_{i_1}, \dots, u_{i_n}}} \mu(\pi, 1_n)$ and

$$W_*^{w_1, \dots, w_n} = \{w \in FP_*^c(G : a) \cup FP_*^c(G : b) : \text{both } L_w^{u_w} \text{ and } L_w^{u_w*} \text{ are in } L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}\}.$$

□

Corollary 3.2. *Let x and y be the D_G -valued random variables in $(W^*(G), E)$. The D_G -valued random variables a and b are free over D_G in $(W^*(G), E)$ if*

$$FP_*(G : P(x, x^*)) \cap FP_*(G : Q(y, y^*)) = \emptyset$$

and

$$W_*^{\{P(x, x^*), Q(y, y^*)\}} = \emptyset,$$

for all $P, Q \in \mathbb{C}[z_1, z_2]$. □

By using (3.2), we can compute the mixed D_G -valued cumulants of two D_G -valued random variables. However, the formula is very abstract. So, we will consider the above formula for fixed two generators of $W^*(G)$.

Definition 3.1. *Let G be a countable directed graph and $\mathbb{F}^+(G)$, the free semi-groupoid of G and let $FP(G)$ be the subset of $\mathbb{F}^+(G)$ consisting of all finite paths. Define a subset $\text{loop}(G)$ of $FP(G)$ containing all loop finite paths or loops. (Remark that, in general, loop finite paths are different from loop-edges. Clearly, all loop-edges are loops in $FP(G)$.) i.e.,*

$$\text{loop}(G) \stackrel{\text{def}}{=} \{l \in FP(G) : l \text{ is a loop}\} \subset FP(G).$$

Also define the subset $\text{loop}^c(G)$ of $FP(G)$ consisting of all non-loop finite path by

$$\text{loop}^c(G) \stackrel{\text{def}}{=} FP(G) \setminus \text{loop}(G).$$

Let $l \in \text{loop}(G)$ be a loop finite path. We say that l is a **basic loop** if there exists no loop $w \in \text{loop}(G)$ such that $l = w^k$, $k \in \mathbb{N} \setminus \{1\}$. Define

$$\text{Loop}(G) \stackrel{\text{def}}{=} \{l \in \text{loop}(G) : l \text{ is a basic loop}\} \subsetneq \text{loop}(G).$$

Let $l_1 = w_1^{k_1}$ and $l_2 = w_2^{k_2}$ in $\text{loop}(G)$, where $w_1, w_2 \in \text{Loop}(G)$. We will say that the loops l_1 and l_2 are **diagram-distinct** if $w_1 \neq w_2$ in $\text{Loop}(G)$. Otherwise, they are not diagram-distinct.

Now, we will introduce the more general diagram-distinctness of general finite paths ;

Definition 3.2. (Diagram-Distinctness) *We will say that the finite paths w_1 and w_2 are **diagram-distinct** if w_1 and w_2 have different diagrams in the graph*

G . Let X_1 and X_2 be subsets of $FP(G)$. The subsets X_1 and X_2 are said to be *diagram-distinct* if x_1 and x_2 are diagram-distinct for all pairs $(x_1, x_2) \in X_1 \times X_2$. This diagram-distinctness implies the diagram-distinctness of loops.

Let H be a directed graph with $V(H) = \{v_1, v_2\}$ and $E(H) = \{e_1 = v_1e_1v_2, e_2 = v_2e_2v_1\}$. Then $l = e_1e_2$ is a loop in $FP(H)$ (i.e., $l \in loop(H)$). Moreover, it is a basic loop (i.e., $l \in Loop(H)$). However, if we have a loop $w = e_1e_2e_1e_2 = l^2$, then it is not a basic loop. i.e.,

$$l^2 \in loop(H) \setminus Loop(H).$$

If the graph G contains at least one basic loop $l \in FP(G)$, then we have

$$\{l^n : n \in \mathbb{N}\} \subseteq loop(G) \quad \text{and} \quad \{l\} \subseteq Loop(G).$$

Suppose that l_1 and l_2 are not diagram-distinct. Then, by definition, there exists $w \in Loop(G)$ such that $l_1 = w^{k_1}$ and $l_2 = w^{k_2}$, for some $k_1, k_2 \in \mathbb{N}$. On the graph G , indeed, l_1 and l_2 make the same diagram. On the other hands, we can see that if $w_1 \neq w_2 \in loop^c(G)$, then they are automatically diagram-distinct.

Lemma 3.3. *Suppose that $w_1 \neq w_2 \in loop^c(G)$ with $w_1 = v_{11}w_1v_{12}$ and $w_2 = v_{21}w_2v_{22}$. Then L_{w_1} and L_{w_2} are free over D_G in $(W^*(G), E)$.*

Proof. By definition, L_{w_1} and L_{w_2} are free over D_G if and only if all mixed D_G -valued cumulants of $W^*(\{L_{w_1}\}, D_G)$ and $W^*(\{L_{w_2}\}, D_G)$ vanish. Equivalently, all D_G -valued cumulants of $P(L_{w_1}, L_{w_1}^*)$ and $Q(L_{w_2}, L_{w_2}^*)$ vanish, for all $P, Q \in \mathbb{C}[z_1, z_2]$. Since $w_1 \neq w_2$ are non-loop edges, we can easily verify that $w_1^{k_1}$ and $w_2^{k_2}$ are not admissible (i.e., $w_1^{k_1} \notin \mathbb{F}^+(G)$ and $w_2^{k_2} \notin \mathbb{F}^+(G)$), for all $k_1, k_2 \in \mathbb{N} \setminus \{1\}$. This shows that

$$L_{w_j}^k = 0_{D_G} = \left(L_{w_j}^k\right)^*, \text{ for } j = 1, 2.$$

Thus, to show that L_{w_1} and L_{w_2} are free over D_G , it suffices to show that all mixed D_G -valued cumulants of $P(L_{w_1}, L_{w_1}^*)$ and $Q(L_{w_2}, L_{w_2}^*)$ vanish, for all $P, Q \in \mathbb{C}[z_1, z_2]$ such that

$$P(z_1, z_2) = \alpha_1 z_1 + \alpha_2 z_1 z_2 + \alpha_3 z_2 z_1 + \alpha_4 z_2$$

and

$$Q(z_1, z_2) = \beta_1 z_1 + \beta_2 z_1 z_2 + \beta_3 z_2 z_1 + \beta_4 z_2,$$

where $\alpha, \beta \in \mathbb{C}$. So, for such P and Q , we have that

$$P(L_{w_1}, L_{w_1}^*) = \alpha_1 L_{w_1} + \alpha_2 L_{v_{11}} + \alpha_3 L_{v_{12}} + \alpha_4 L_{w_1}^*$$

and

$$Q(L_{w_2}, L_{w_2}^*) = \beta_1 L_{w_2} + \beta_2 L_{v_{21}} + \beta_3 L_{v_{22}} + \beta_4 L_{w_2}^*.$$

Thus, we have that

$$FP_*(G : P(L_{w_1}, L_{w_1}^*)) \supseteq \{w_1\}, \quad FP_*(G : Q(L_{w_2}, L_{w_2}^*)) \supseteq \{w_2\}$$

and

$$FP_*^c(G : P(L_{w_1}, L_{w_1}^*)) \supseteq \{w_1\}, \quad FP_*^c(G : Q(L_{w_2}, L_{w_2}^*)) \supseteq \{w_2\}.$$

Remark that if $FP_*(G : P(L_{w_1}, L_{w_1}^*)) = \{w_1\}$, then $FP_*^c(G : P(L_{w_1}, L_{w_1}^*)) = \emptyset$, and if $FP_*^c(G : P(L_{w_1}, L_{w_1}^*)) = \{w_1\}$, then $FP_*(G : P(L_{w_1}, L_{w_1}^*)) = \emptyset$. The similar relation holds for $Q(L_{w_2}, L_{w_2}^*)$. So, we have that

$$FP_*(G : P(L_{w_1}, L_{w_1}^*)) \cap FP_*(G : Q(L_{w_2}, L_{w_2}^*)) = \emptyset$$

and

$$W_*^{\{P(L_{w_1}, L_{w_1}^*), Q(L_{w_2}, L_{w_2}^*)\}} = \emptyset.$$

Therefore, by the formula (3.4.3), we have the vanishing mixed D_G -valued cumulants of $P(L_{w_1}, L_{w_1}^*)$ and $Q(L_{w_2}, L_{w_2}^*)$, for all $n \in \mathbb{N}$ and for all such $P, Q \in \mathbb{C}[z_1, z_2]$. So, we can conclude that L_{w_1} and L_{w_2} are free over D_G in $(W^*(G), E)$. ■

Now, we will consider the loop case.

Lemma 3.4. *Let $l_1 \neq l_2 \in \text{Loop}(G)$ be **basic loops** such that $l_1 = v_1 l_1 v_1$ and $l_2 = v_2 l_2 v_2$, for $v_1, v_2 \in V(G)$ (possibly $v_1 = v_2$). i.e.e, two basic loops l_1 and l_2 are diagram-distinct. Then the D_G -valued random variables L_{l_1} and L_{l_2} are free over D_G in $(W^*(G), E)$.*

Proof. Different from the non-loop case, if l_1 and l_2 are loops, then $l_1^{k_1}$ and $l_2^{k_2}$ exist in $FP(G)$, for all $k_1, k_2 \in \mathbb{N}$. To show that L_{l_1} and L_{l_2} are free over D_G , it suffices to show that all mixed D_G -valued cumulants of $P(L_{w_1}, L_{w_1}^*)$ and $Q(L_{w_2}, L_{w_2}^*)$ vanish, for all $P, Q \in \mathbb{C}[z_1, z_2]$. such that

$$P(z_1, z_2) = f_1(z_1) + f_2(z_2) + P_0(z_1, z_2)$$

and

$$Q(z_1, z_2) = g_1(z_1) + g_2(z_2) + Q_0(z_1, z_2),$$

where $f_1, f_2, g_1, g_2 \in \mathbb{C}[z]$ and $P_0, Q_0 \in \mathbb{C}[z_1, z_2]$ such that P_0 and Q_0 does not contain polynomials only in z_1 and z_2 . So, for such P and Q , we have that

$$P(L_{l_1}, L_{l_1}^*) = f_1(L_{l_1}) + f_2(L_{l_1}^*) + P_0(L_{l_1}, L_{l_1}^*)$$

and

$$Q(L_{l_2}, L_{l_2}^*) = g_1(L_{l_2}) + g_2(L_{l_2}^*) + Q_0(L_{l_2}, L_{l_2}^*).$$

Notice that $L_{l_j}^k = L_{l_j^k}$, for all $k \in \mathbb{N}$, $j = 1, 2$. Also, notice that

$$P_0(L_{l_1}, L_{l_1}^*) = f_1^0(L_{l_1}) + f_2^0(L_{l_1}^*) + \alpha L_{v_1}$$

and

$$Q_0(L_{l_2}, L_{l_2}^*) = g_1^0(L_{l_2}) + g_2^0(L_{l_2}^*) + \beta L_{v_2},$$

where $f_1^0, f_2^0, g_1^0, g_2^0 \in \mathbb{C}[z]$ and $\alpha, \beta \in \mathbb{C}$, by the fact that

$$L_{l_j}^* L_{l_j} = L_{v_j} = L_{l_j} L_{l_j}^*,$$

under the weak-topology. So, finally, we have that

$$P(L_{l_1}, L_{l_1}^*) = \mathbf{f}_1(L_{l_1}) + \mathbf{f}_2(L_{l_1}^*) + \alpha L_{v_1}$$

and

$$Q(L_{l_2}, L_{l_2}^*) = \mathbf{g}_1(L_{l_2}) + \mathbf{g}_2(L_{l_2}^*) + \beta L_{v_2},$$

where $\mathbf{f}_1, \mathbf{f}_2, \mathbf{g}_1, \mathbf{g}_2 \in \mathbb{C}[z]$ and $\alpha, \beta \in \mathbb{C}$. Thus, we have that

$$FP_*(G : P(L_{l_1}, L_{l_1}^*)) \subseteq \{l_1^k\}_{k=1}^\infty, \quad FP_*(G : Q(L_{l_2}, L_{l_2}^*)) \subseteq \{l_2^k\}_{k=1}^\infty$$

and

$$FP_*^c(G : P(L_{l_1}, L_{l_1}^*)) \subseteq \{l_1^k\}_{k=1}^\infty, \quad FP_*^c(G : Q(L_{l_2}, L_{l_2}^*)) \subseteq \{l_2^k\}_{k=1}^\infty.$$

So, we have that

$$FP_*(G : P(L_{w_1}, L_{w_1}^*)) \cap FP_*(G : Q(L_{w_2}, L_{w_2}^*)) = \emptyset,$$

because l_1 and l_2 are in $Loop(G)$ (and hence if $l_1 \neq l_2$, then they are diagram-distinct.) And we have that

$$W_*^{\{P(L_{w_1}, L_{w_1}^*), Q(L_{w_2}, L_{w_2}^*)\}} = \emptyset.$$

Therefore, by the formula (3.4.3), we have the vanishing mixed D_G -valued cumulants of $P(L_{l_1}, L_{l_1}^*)$ and $Q(L_{l_2}, L_{l_2}^*)$, for all $n \in \mathbb{N}$ and for all $P, Q \in \mathbb{C}[z_1, z_2]$. Since P and Q are arbitrary, we can conclude that L_{l_1} and L_{l_2} are free over D_G in $(W^*(G), E)$. ■

Notice that we assumed that the loops l_1 and l_2 are basic loops in the previous lemma. Since they are distinct basic loops, they are automatically diagram-distinct. Now, assume that l_1 and l_2 are not diagram-distinct. i.e., there exists a basic loop $w \in Loop(G)$ such that $l_1 = w^{k_1}$ and $l_2 = w^{k_2}$, for some $k_1, k_2 \in \mathbb{N}$. In other words, the loops l_1 and l_2 have the same diagram in the graph G . Then the D_G -valued random variables L_{l_1} and L_{l_2} are not free over D_G in $(W^*(G), E)$. See the next example ;

Example 3.1. Let G_1 be a directed graph with $V(G_1) = \{v\}$ and $E(G_1) = \{l = vl\}$. So, in this case,

$$\begin{aligned} E(G_1) &= \text{Loop}(G_1), \quad FP(G_1) = \text{loop}(G_1), \\ \text{and} \quad \text{loop}(G_1) &= \{l^k : k \in \mathbb{N}\}. \end{aligned}$$

Thus, even if $w_1 \neq w_2 \in \text{loop}(G_1)$, w_1 and w_2 are Not diagram-distinct. Take l^2 and l^3 in $FP(G_1)$. Then the D_{G_1} -valued random variable L_{l^2} and L_{l^3} are not free over D_{G_1} in $(W^*(G_1), E)$. Indeed, let's take $P, Q \in \mathbb{C}[z_1, z_2]$ as

$$P(z_1, z_2) = z_1^3 + z_2^3 \quad \text{and} \quad Q(z_1, z_2) = z_1^2 + z_2^2.$$

Then

$$P(L_{l^2}, L_{l^2}^*) = L_{l^2}^3 + L_{l^2}^{*3} = L_{l^6} + L_{l^6}^*$$

and

$$Q(L_{l^3}, L_{l^3}^*) = L_{l^3}^2 + L_{l^3}^{*2} = L_{l^6} + L_{l^6}^*.$$

Then

$$\begin{aligned} k_2(P(L_{l^2}, L_{l^2}^*), Q(L_{l^3}, L_{l^3}^*)) &= k_2(L_{l^6} + L_{l^6}^*, L_{l^6} + L_{l^6}^*) \\ &= \mu_{l^6, l^6}^{1,*} \text{Pr} \text{oj}(L_{l^6}, L_{l^6}^*) + \mu_{l^6, l^6}^{*,1} \text{Pr} \text{oj}(L_{l^6}^*, L_{l^6}) \\ &= \mu_{l^6, l^6}^{1,*} L_v + \mu_{l^6, l^6}^{*,1} L_v = (\mu_{l^6, l^6}^{1,*} + \mu_{l^6, l^6}^{*,1}) L_v \\ &= 2L_v \neq 0_{D_G}, \end{aligned}$$

since $\mu_{l^6, l^6}^{1,*} = \mu(1_2, 1_2) = 1 = \mu_{l^6, l^6}^{*,1}$. This says that there exists at least one nonvanishing mixed D_G -valued cumulant of $W^*(\{L_{l^3}\}, D_{G_1})$ and $W^*(\{L_{l^2}\}, D_{G_1})$. Therefore, L_{l^3} and L_{l^2} are not free over D_{G_1} in $(W^*(G_1), E)$.

As we have seen before, if two loops l_1 and l_2 are not diagram-distinct, then D_G -valued random variables L_{l_1} and L_{l_2} are Not free over D_G . However, if l_1 and l_2 are diagram-distinct, we can have the following lemma, by the previous lemma ;

Lemma 3.5. Let $l_1 \neq l_2 \in \text{loop}(G)$ be loops and assume that $l_1 = w_1^{k_1}$ and $l_2 = w_2^{k_2}$, where $w_1, w_2 \in \text{Loop}(G)$ are basic loops and $k_1, k_2 \in \mathbb{N}$. If $w_1 \neq w_2 \in \text{Loop}(G)$, then the D_G -valued random variables L_{l_1} and L_{l_2} are free over D_G in $(W^*(G), E)$. $\square$

Finally, we will observe the following case when we have a loop and a non-loop finite path.

Lemma 3.6. Let $l \in \text{loop}(G)$ and $w \in \text{loop}^c(G)$. Then the D_G -valued random variables L_l and L_w are free over D_G in $(W^*(G), E)$.

Proof. let $l \in \text{loop}(G)$ and $w \in \text{loop}^c(G)$ and let L_l and L_w be the corresponding D_G -valued random variables in $(W^*(G), E)$. Then, for all $P, Q \in \mathbb{C}[z_1, z_2]$, we have that

$$FP_*(G : P(L_l, L_l^*)) \cap FP_*(G : Q(L_w, L_w^*)) = \emptyset,$$

since

$$FP_*(G : P(L_l, L_l^*)) \subseteq \{l^k : k \in \mathbb{N}\} \subset \text{loop}(G)$$

and

$$FP_*(G : Q(L_w, L_w^*)) = \{w\} \subset \text{loop}^c(G).$$

Also, since $\text{loop}(G) \cap \text{loop}^c(G) = \emptyset$, we can get that

$$W_*^{\{P(L_l, L_l^*), Q(L_w, L_w^*)\}} = \emptyset,$$

for all $P, Q \in \mathbb{C}[z_1, z_2]$. Therefore, the D_G -valued random variables L_l and L_w are free over D_G in $(W^*(G), E)$. ■

Now, we can summarize the above lemmas in this section as follows and this theorem is one of the main result of this paper. The theorem is the characterization of D_G -freeness of generators of $W^*(G)$ over D_G .

Theorem 3.7. *Let $w_1, w_2 \in FP(G)$ be finite paths. The D_G -valued random variables L_{w_1} and L_{w_2} in $(W^*(G), E)$ are free over D_G if and only if w_1 and w_2 are diagram-distinct.*

Proof. $\Leftarrow$) Suppose that finite paths w_1 and w_2 are diagram-distinct. Then the D_G -valued random variables L_{w_1} and L_{w_2} are free over D_G , by the previous lemmas.

$\Rightarrow$) Let L_{w_1} and L_{w_2} be free over D_G in $(W^*(G), E)$. Now, assume that w_1 and w_2 are not diagram-distinct. We will observe the following cases ;

(Case I) The finite paths $w_1, w_2 \in \text{loop}(G)$. Since they are not diagram-distinct, there exists a basic loop $l = vlv \in \text{Loop}(G)$, with $v \in V(G)$, such that $w_1 = l^{k_1}$ and $w_2 = l^{k_2}$, for some $k_1, k_2 \in \mathbb{N}$. As we have seen before, L_{w_1} and L_{w_2} are not free over D_G in $(W^*(G), E)$. Indeed, if we let $k \in \mathbb{N}$ such that $k_1 \mid k$ and $k_2 \mid k$ with $k = k_1 n_1 = k_2 n_2$, for $n_1, n_2 \in \mathbb{N}$, then we can take $P, Q \in \mathbb{C}[z_1, z_2]$ defined by

$$P(z_1, z_2) = z_1^{n_1} + z_2^{n_1} \quad \text{and} \quad Q(z_1, z_2) = z_1^{n_2} + z_2^{n_2}.$$

And then

$$P(L_{w_1}, L_{w_1}^*) = L_{w_1}^{n_1} + L_{w_1}^{*n_1} = L_{l^{k_1}}^{n_1} + L_{l^{k_1}}^{*n_1} = L_{l^k} + L_{l^k}^*$$

and

$$Q(L_{w_2}, L_{w_2}^*) = L_{w_2}^{n_2} + L_{w_2}^{*n_2} = L_{l^{k_2}}^{n_2} + L_{l^{k_2}}^{*n_2} = L_{l^k} + L_{l^k}^*.$$

So,

$$\begin{aligned} k_2(P(L_{w_1}, L_{w_1}^*), Q(L_{w_2}, L_{w_2}^*)) &= k_2(L_{l^k} + L_{l^k}^*, L_{l^k} + L_{l^k}^*) \\ &= \mu_{l^k, l^k}^{1,*} L_v + \mu_{l^k, l^k}^{*,1} L_v \\ &= 2L_v \neq 0_{D_G}. \end{aligned}$$

Therefore, $P(L_{w_1}, L_{w_1}^*)$ and $Q(L_{w_2}, L_{w_2}^*)$ are not free over D_G . This shows that $W^*(\{L_{w_1}\}, D_G)$ and $W^*(\{L_{w_2}\}, D_G)$ are not free over D_G in $(W^*(G), E)$ and hence L_{w_1} and L_{w_2} are not free over D_G . This contradict our assumption.

(Case II) Suppose that the finite paths w_1, w_2 are non-loop finite paths in $\text{loop}^c(G)$ and assume that they are not diagram-distinct. Since they are not diagram-distinct, they are identically equal. Therefore, they are not free over D_G in $(W^*(G), E)$.

(Case III) Let $w_1 \in \text{loop}(G)$ and $w_2 \in \text{loop}^c(G)$. They are always diagram-distinct.

Let L_{w_1} and L_{w_2} are free over D_G and assume that w_1 and w_2 are not diagram-distinct. Then L_{w_1} and L_{w_2} are not free over D_G , by the Case I, II and III. So, this contradict our assumption. ■

The previous theorem characterize the D_G -freeness of two partial isometries L_{w_1} and L_{w_2} , where $w_1, w_2 \in FP(G)$. This characterization shows us that the diagram-distinctness of finite paths determine the D_G -freeness of corresponding creation operators.

Let a and b be the given D_G -valued random variables in (3.0). We can get the necessary condition for the D_G -freeness of a and b , in terms of their supports. Recall that we say that the two subsets X_1 and X_2 of $FP(G)$ are said to be diagram-distinct if x_1 and x_2 are diagram-distinct, for all pairs $(x_1, x_2) \in X_1 \times X_2$.

Theorem 3.8. *Let $a, b \in (W^*(G), E)$ be D_G -valued random variables with their supports $\mathbb{F}^+(G : a)$ and $\mathbb{F}^+(G : b)$. The D_G -valued random variables a and b are free over D_G in $(W^*(G), E)$ if $FP(G : a_1)$ and $FP(G : a_2)$ are diagram-distinct.*

Proof. For convenience, let's denote a and b by a_1 and a_2 , respectively. Assume that the supports of a_1 and a_2 , $\mathbb{F}^+(G : a_1)$ and $\mathbb{F}^+(G : a_2)$ are diagram-distinct. Then by the previous D_G -freeness characterization,

$$\sum_{l \in FP(G:a_1), u_l \in \{1,*\}} p_l^{(1)} L_l^{u_l} \quad \text{and} \quad \sum_{l \in FP(G:a_2), u_l \in \{1,*\}} p_l^{(2)} L_l^{u_l}$$

are free over D_G in $(W^*(G), E)$. Indeed, since $FP(G : a_1)$ and $FP(G : a_2)$ are diagram-distinct, all summands L_{w_1} 's of a_1 and L_{w_2} 's of a_2 are free over D_G in $(W^*(G), E)$. Therefore, a_1 and a_2 are free over D_G in $(W^*(G), E)$. ■

4. D_G -VALUED SEMICIRCULAR ELEMENTS

Throughout this chapter, we will consider the D_G -valued semicircularity. Let B be a von Neumann algebra and A , a W^* -algebra over B and let $E : A \rightarrow B$ be a conditional expectation. Then (A, E) be an amalgamated W^* -probability space over B . We say that the D_G -valued random variable $x \in (A, E)$ is a B -valued semicircular element if it is self-adjoint and the only nonvanishing B -valued cumulants is the second one. i.e., $x \in (A, E)$ is B -semicircular if x is self-adjoint in A and

$$k_n(x, \dots, x) = \begin{cases} k_2(x, x) \neq 0_B & \text{if } n = 2 \\ 0_B & \text{otherwise.} \end{cases}$$

Let G be a directed graph with $V(G) = \{v\}$ and $E(G) = \{e = vev\}$. i.e., G is a one-vertex graph with one loop edge. Canonically, we can construct the graph W^* -algebra $W^*(G)$ and its diagonal subalgebra $D_G \simeq \Delta_1 = \mathbb{C}$. So, the canonical conditional expectation E is a linear functional. Moreover, it is a trace on $W^*(G)$. Notice that, by Voiculescu,

$$W^*(G) \simeq L(\mathbb{Z}),$$

where $L(\mathbb{Z})$ is the free group factor. Moreover, the random variable $L_e + L_e^*$ is the Voiculescu's semicircular element. Therefore, the only nonvanishing cumulants of $L_e + L_e^*$ is the second one. First, we will consider the following combinatorial fact. This is crucial to consider the D_G -semicircularity on $(W^*(G), E)$.

Lemma 4.1. *Let G and e be given as above. Then*

$$\sum_{L \in LP_2^*} \mu_{e,e}^{L(u_1, u_2)} = 2$$

and

$$\sum_{L \in LP_n^*} \mu_{e, \dots, e}^{L(u_1, \dots, u_n)} = 0, \quad \forall n \in 2\mathbb{N} \setminus \{2\}.$$

Proof. Define $a = L_e + L_e^*$. Then it is a semicircular element, in the sense of Voiculescu. So, the only nonvanishing D_G -valued cumulant of a is the second one. By the previous lemma, we have that

$$k_n(a, \dots, a) = \sum_{L \in LP_n^*} \mu_{e, \dots, e}^{L(u_1, \dots, u_n)} \cdot L_v.$$

Suppose that $n = 2$. Then

$$k_2(a, a) = \mu_{e, e}^{1, *} \cdot L_v + \mu_{e, e}^{*, 1} \cdot L_v = \mu_{e, e}^{1, *} + \mu_{e, e}^{*, 1},$$

since $L_v = 1_{D_G} = 1_{\mathbb{C}} = 1$. Notice that $l_e l_{e^{-1}}$ and $l_{e^{-1}} l_e$ have their lattice paths

$$* \nearrow \searrow \quad \text{and} \quad * \searrow \nearrow,$$

respectively. Therefore, $C_{e, e}^{1, *} = C_{e, e}^{*, 1} = \{1_2\}$. Therefore,

$$\mu_{e, e}^{1, *} = \mu_{e, e}^{*, 1} = \mu(1_2, 1_2) = 1.$$

Therefore, $k_2(a, a) = 2$. Equivalently,

$$\sum_{L \in LP_2^*} \mu_{e, e}^{L(u_1, u_2)} = 2,$$

since $L_v = 1_{D_G} = 1_{\mathbb{C}} = 1$. Now, let $2 < n \in 2\mathbb{N}$. Then

$$k_n(a, \dots, a) = \sum_{L \in LP_n^*} \mu_{e, \dots, e}^{L(u_1, \dots, u_n)} \cdot L_v = \sum_{L \in LP_n^*} \mu_{e, \dots, e}^{L(u_1, \dots, u_n)}$$

since $L_v = 1$

$$= 0,$$

by the semicircularity of $L_e + L_e^*$. Therefore,

$$\sum_{L \in LP_n^*} \mu_{e, \dots, e}^{L(u_1, \dots, u_n)} = 0, \quad \forall n \in 2\mathbb{N} \setminus \{2\}. \blacksquare$$

By the previous lemma, we can determine the D_G -semicircular elements in $(W^*(G), E)$.

Theorem 4.2. *Let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over the diagonal subalgebra D_G . Let $w = vwv \in \text{loop}(G)$, with $v \in V(G)$. Then $L_w + L_w^*$ is a D_G -valued semicircular elements, with*

$$k_2(L_w + L_w^*, L_w + L_w^*) = 2L_v.$$

Proof. Let $w = v w v \in \text{loop}(G)$, with $v \in V(G)$. Define a D_G -valued random variable $a = L_w + L_w^*$. Then, clearly, it is self-adjoint in $W^*(G)$. It suffices to show that it has only second nonvanishing D_G -valued cumulants.

$$\begin{aligned} k_n(a, \dots, a) &= k_n(L_w + L_w^*, \dots, L_w + L_w^*) \\ &= \sum_{(u_1, \dots, u_n) \in \{1, *\}^n} k_n(L_w^{u_1}, \dots, L_w^{u_n}) \end{aligned}$$

by the bimodule map property of k_n

$$= \sum_{l_{w^{t_1}} \dots l_{w^{t_n}} : * \text{-axis-property}} k_n(L_w^{u_1}, \dots, L_w^{u_n})$$

where $l_{w^{t_1}} \dots l_{w^{t_n}} \in (A_{G^*} / R, E)$ is the corresponding element of $L_w^{u_1} \dots L_w^{u_n}$

$$= \sum_{L \in LP_n^*} \mu_{w, \dots, w}^{L(u_1, \dots, u_n)} \cdot L_v.$$

But by the previous theorem,

$$\sum_{L \in LP_n^*} \mu_{w, \dots, w}^{L(u_1, \dots, u_n)} = \begin{cases} 2 & \text{if } n = 2 \\ 0 & \text{otherwise.} \end{cases}$$

Therefore,

$$k_n(a, \dots, a) = \begin{cases} 2L_v & \text{if } n = 2 \\ 0_{D_G} & \text{otherwise.} \end{cases}$$

and hence $a = L_w + L_w^*$, $w \in \text{loop}(G)$, is a D_G -semicircular element in $(W^*(G), E)$. Notice that if w is a loop (as a finite path), then $C_{w, \dots, w}^{L(u_1, \dots, u_n)} = C_{e, \dots, e}^{L(u_1, \dots, u_n)}$, where e is the given at the beginning of this chapter. Therefore, we have the previous formul. ■

So, we can conclude that all D_G -valued random variables having the forms of $pL_w + pL_w^*$ ($p \in \mathbb{R}$, $w \in \text{loop}(G)$) are D_G -semicircular elements in $(W^*(G), E)$.

Corollary 4.3. *Let $w_j = v_j w_j v_j \in \text{loop}(G)$, $v_j \in V(G)$, $j = 1, \dots, N$, such that they are **mutually diagram-distinct** (Note that it is possible that $v_i = v_j$, for any $i \neq j$ in $\{1, \dots, N\}$). Then the D_G -valued random variable*

$$a = \sum_{j=1}^N (p_j L_{w_j} + p_j L_{w_j}^*) \in (W^*(G), E)$$

is D_G -semicircular and

$$k_2(a, a) = \sum_{j=1}^N 2p_j^4 \cdot L_{v_j}.$$

□

5. D_G -EVEN ELEMENTS

Let G be a directed graph with $V(G) = \{v_1, v_2\}$ and $E(G) = \{e = v_1 e v_2\}$. We can construct the graph W^* -algebra $W^*(G)$ and its diagonal subalgebra $D_G = \Delta_2$. Trivially $L_{v_1} + L_{v_2} = 1_{D_G}$. Define a D_G -valued random variable $a = L_e + L_e^*$. Note that

$$FP(G) = E(G),$$

since $e^k \notin \mathbb{F}^+(G)$, for all $k \in \mathbb{N} \setminus \{1\}$. We cannot construct the more finite paths other than e , itself. Also, the support of this operator a is

$$\mathbb{F}^+(G : a) = FP(G : a) = FP_*(G : a) = E(G).$$

Similar to the previous chapter, we will observe the following combinatorial fact ;

Lemma 5.1. *Let G and e be given as above. Then*

$$\sum_{L \in LP_n^*} \mu_{e, \dots, e}^{L(u_1, \dots, u_n)} = 2\mu_n \in \mathbb{R},$$

where $\mu_n = \mu_{e, e, \dots, e, e}^{1, *, \dots, 1, *} = \mu_{e, e, \dots, e, e}^{*, 1, \dots, *, 1}$, for all $n \in 2\mathbb{N}$.

Proof. Since $e = v_1 e v_2$, with $v_1 \neq v_2 \in V(G)$, $e^k \notin \mathbb{F}^+(G)$ (i.e., it is not admissible), whenever $n > 1$. Therefore, for any even n ,

$$\sum_{L \in LP_n^*} \mu_{e, \dots, e}^{L(u_1, \dots, u_n)} = \mu_{e, e, \dots, e, e}^{1, *, \dots, 1, *} + \mu_{e, e, \dots, e, e}^{*, 1, \dots, *, 1}.$$

Indeed, if we let $(u_1, \dots, u_n) \in \{1, *\}^n$ and if it is not alternating, then there exists at least one $j \in \{1, \dots, n\}$ such that $u_j = u_{j+1}$ in $\{1, *\}$. This means that there should be a consecutive increasing or decreasing words of e 's. But e^k does not exist in our graph G , whenever $k > 1$. Therefore, for such $L(u_1, \dots, u_n) \in LP_n^*$,

$$\mu_{e, \dots, e}^{L(u_1, \dots, u_n)} = \sum_{\theta \in C_{e, \dots, e}^{L(u_1, \dots, u_n)}} \mu(\theta, 1_n) = 0 \in \mathbb{R}.$$

Observe that $C_{e,e,\dots,e,e}^{1,*,\dots,1,*}$ and $C_{e,e,\dots,e,e}^{*,1,\dots,*,1}$ have the same elements, by the symmetry of the lattice paths of $l_e l_{e-1} \dots l_e l_{e-1}$ and $l_{e-1} l_e \dots l_{e-1} l_e$. So,

$$\mu_{e,e,\dots,e,e}^{1,*,\dots,1,*} = \mu_{e,e,\dots,e,e}^{*,1,\dots,*,1}.$$

Denote this real value by μ_n , for each $n \in 2\mathbb{N}$. Then we can get the above formula. ■

So, we have that ;

Lemma 5.2. *Let G and e be given as above. Then the D_G -valued cumulants of $a = L_e + L_e^*$ is determined by ;*

(1) $k_n(a, \dots, a) = 0_{D_G}$, whenever n is odd.

(2) $k_n(a, \dots, a) = \mu_n \cdot 1_{D_G}$, for all $n \in 2\mathbb{N}$, where

$$\mu_n = \mu_{e,e,\dots,e,e}^{1,*,\dots,1,*} = \mu_{e,e,\dots,e,e}^{*,1,\dots,*,1}.$$

Proof. Suppose n is odd. Then there is no lattice path having the $*$ -axis-property. Therefore, (1) holds true. Now, assume that $n \in 2\mathbb{N}$. Let $L \in LP_n^*$. Since e is a non-loop edge, $e^k \notin FP(G) = E(G)$. Therefore, when we consider $l_{e^{t_1}} \dots l_{e^{t_n}} \in (A_G^- / R, E)$, the corresponding element of $L_e^{u_1} \dots L_e^{u_n}$ (where L_e and L_e^* are summands of a), the lattice paths of it is

$$* \nearrow \searrow \nearrow \searrow \dots \nearrow \searrow \quad \text{or} \quad * \searrow \nearrow \searrow \nearrow \dots \searrow \nearrow,$$

denoted by $[rfrf \dots rfr]$ and $[frfr \dots fr]$, respectively (Here r stands for the rising step and f stands for the falling step), because if there are consecutive rising steps or consecutive falling steps, then it represents $l_e \dots l_e = 0_{D_G}$ or $l_{e-1} \dots l_{e-1}$. Therefore, if $n \in 2\mathbb{N}$, then

$$\begin{aligned} & k_n(a, \dots, a) \\ &= \sum_{(u_1, \dots, u_n) \in \{1, *\}^n} k_n(L_e^{u_1}, \dots, L_e^{u_n}) \\ &= \sum_{L \in LP_n^*} \mu_{e,e,\dots,e}^{L(u_1, \dots, u_n)} \cdot E(L_e^{u_1} \dots L_e^{u_n}) \\ &= \mu_{e,e,\dots,e,e}^{1,*,\dots,1,*} E(L_e L_e^* \dots L_e L_e^*) + \mu_{e,e,\dots,e,e}^{*,1,\dots,*,1} E(L_e^* L_e \dots L_e^* L_e) \\ &= \mu_n \cdot L_{v_1} + \mu_n L_{v_2} = \mu_n (L_{v_1} + L_{v_2}) \end{aligned}$$

where $\mu_n = \mu_{e,e,\dots,e,e}^{1,*,\dots,1,*} = \mu_{e,e,\dots,e,e}^{*,1,\dots,*,1}$

$$= \mu_n \cdot 1_{D_G}.$$

Indeed, the lattice paths $[rf...rf]$ and $[fr...fr]$ in LP_n^* induce the same sets $C_{e,e,\dots,e,e}^{1,*,\dots,1,*}$ and $C_{e,e,\dots,e,e}^{*,1,\dots,*,1}$. Thus

$$\mu_{e,e,\dots,e,e}^{1,*,\dots,1,*} = \sum_{\theta \in C_{e,e,\dots,e,e}^{1,*,\dots,1,*}} \mu(\theta, 1_n) = \sum_{\pi \in C_{e,e,\dots,e,e}^{*,1,\dots,*,1}} \mu(\pi, 1_n) = \mu_{e,e,\dots,e,e}^{*,1,\dots,*,1}.$$

■

Now, we will observe the general case. Define $\mu_n \stackrel{\text{def}}{=} \mu_{e,e,\dots,e,e}^{1,*,\dots,1,*} = \mu_{e,e,\dots,e,e}^{*,1,\dots,*,1}$,

where e is given as before. Now, we will observe the general case when e is a general finite path in an arbitrary countable directed graph G .

Definition 5.1. Let B be a von Neumann algebra and let A be the W^* -algebra over B . Let $E_B : A \rightarrow B$ be a conditional expectation and (A, E_B) , the amalgamated W^* -probability space over B . We say that the B -valued random variable $a \in (A, E_B)$ is B -valued even (in short, B -even) if it is self-adjoint and if it has all vanishing odd B -valued moments.

Let B be a von Neumann algebra and A , a W^* -algebra over B and let (A, E_B) be the amalgamated W^* -probability space over B . Recall that B -valued random variable $a \in (A, E_B)_{sa}$ is B -even if and only if all odd B -valued cumulants of a vanish. By the previous lemma, we can easily see that D_G -valued random variables $L_w + L_w^*$, for $w \in FP(G)$, are D_G -even, because if $w = v_1 w v_2$, then

$$k_n(L_w + L_w^*, \dots, L_w + L_w^*) = \begin{cases} 0_{D_G} & \text{if } n \text{ is odd} \\ \mu_n \cdot (L_{v_1} + L_{v_2}), & \text{if } n \text{ is even} \end{cases}$$

for all $n \in \mathbb{N}$, where $\mu_n = \mu_{w,w,\dots,w,w}^{1,*,\dots,1,*} = \mu_{w,w,\dots,w,w}^{*,1,\dots,*,1}$. Remark that

$$C_{w,w,\dots,w,w}^{1,*,\dots,1,*} = C_{w,w,\dots,w,w}^{*,1,\dots,*,1} = C_{e,e,\dots,e,e}^{*,1,\dots,*,1} = C_{e,e,\dots,e,e}^{1,*,\dots,1,*},$$

where e is the edge given at the beginning of this chapter. Based on it, we can get that

Theorem 5.3. Let $w \in FP(G)$. Then the D_G -valued random variable $L_w + L_w^*$ is D_G -even. Moreover, if $w = v_1 w v_2$ with $v_1, v_2 \in V(G)$ (possibly $v_1 = v_2$), then

$$k_n(L_w + L_w^*, \dots, L_w + L_w^*) = \begin{cases} 0_{D_G} & \text{if } n \text{ is odd} \\ \mu_n \cdot (L_{v_1} + L_{v_2}), & \text{if } n \text{ is even} \end{cases}$$

for all $n \in \mathbb{N}$, where $\mu_n = \mu_{w,w,\dots,w,w}^{1,*,\dots,1,*} = \mu_{w,w,\dots,w,w}^{*,1,\dots,*,1}$. □

6. D_G -VALUED R-DIAGONAL ELEMENTS

In this chapter, we will consider the D_G -valued R-diagonality on the graph W^* -probability space $(W^*(G), E)$ over the diagonal subalgebra D_G . Recall that

Definition 6.1. *Let B be a von Neumann algebra and let A be a W^* -algebra over B . Suppose that we have a conditional expectation $E_B : A \rightarrow B$ and hence (A, E_B) is the amalgamated W^* -probability space over B . We say that the D_G -valued random variable $x \in (A, E_B)$ is a D_G -valued R-diagonal element if the only nonvanishing mixed cumulants of x and x^* are*

$$\begin{aligned} & k_n(b_1x, b_2x^*, \dots, b_{n-1}x, b_nx^*) \\ \text{and} \\ & k_n(b'_1x^*, b_2x, \dots, b'_{n-1}x^*, b'_nx), \end{aligned}$$

for all $n \in 2\mathbb{N}$, where $b_1, b'_1, \dots, b_n, b'_n \in B$ are arbitrary. If n is odd, then automatically the mixed cumulants vanish.

We can show that L_w is D_G -valued R-diagonal, whenever w is a finite path in G . By the results in the proceeding two chapters, we can get the following theorem ;

Theorem 6.1. *Let G be a countable directed graph and $w \in FP(G)$. Then the D_G -valued random variable L_w and $L_w^* \in (W^*(G), E)$ are D_G -valued R-diagonal.*

Proof. It suffices to show that the only nonvanishing mixed cumulants of L_w and L_w^* are alternating ones. i.e., the nonvanishing mixed cumulants are

$$(6.1) \quad k_{2n}(L_w^*, L_w, \dots, L_w^*, L_w) \quad \text{and} \quad k_{2n}(L_w, L_w^*, \dots, L_w, L_w^*).$$

Suppose that w is a loop. Then by Lemma 4.1, we have that the only nonvanishing mixed cumulants are

$$k_2(L_w^*, L_w) \quad \text{and} \quad k_2(L_w, L_w^*).$$

So, L_w and L_w^* are D_G -valued R-diagonal.

Now, assume that $w = v_1 w v_2$ is a non-loop finite path, with $v_1 \neq v_2 \in V(G)$. Then the nonvanishing mixed cumulants of L_w and L_w^* have the forms of (6.1). By Section 2.2, we can easily get that

$$k_{2n}(L_w^*, L_w, \dots, L_w^*, L_w) = \mu_{w, w, \dots, w, w}^{*, 1, \dots, *, 1} \cdot L_{v_1}$$

and

$$k_{2n}(L_w, L_w^*, \dots, L_w, L_w^*) = \mu_{w, w, \dots, w, w}^{1, *, \dots, 1, *} \cdot L_{v_2}.$$

Assume that there exists a nonvanishing mixed D_G -cumulant of L_w and L_w^* . i.e., assume that there exist $n \in \mathbb{N}$ and a n -tuple $(u_1, \dots, u_n)$ of $\{1, *\}$ such that $k_n(L_w^{u_1}, \dots, L_w^{u_n}) \neq 0_{D_G}$. By Section 2.2, we have that

$$k_n(L_w^{u_1}, \dots, L_w^{u_n}) = \mu_{w, \dots, w}^{u_1, \dots, u_n} E(L_w^{u_1} \dots L_w^{u_n}).$$

Notice that since w is a non-loop finite path, there is no admissible finite path w^k , for $k \in \mathbb{N} \setminus \{1\}$. So, if $(u_1, \dots, u_n)$ is not alternating, then there exists at least one j in $\{1, \dots, n-1\}$ such that $u_j = u_{j+1}$. Since $L_w^{u_j} L_w^{u_{j+1}} = L_w^2$ or L_w^{*2} , the D_G -valued random variable $L_w^{u_1} \dots L_w^{u_n}$ does not have the $*$ -axis-property and hence

$$E(L_w^{u_1} \dots L_w^{u_n}) = 0_{D_G} = k_n(L_w^{u_1}, \dots, L_w^{u_n}).$$

This contradict our assumption. So, L_w and L_w^* are D_G -valued R-diagonal. ■

The above theorem shows us that all generators of $W^*(G)$ generated by finite paths in $FP(G)$ are D_G -valued R-diagonal.

7. GENERATING OPERATORS

In this chapter, as examples, we will compute the trivial D_G -valued moments and cumulants of the generating operator T of the graph W^* -algebra $W^*(G)$. Let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over its diagonal subalgebra. Let $a \in (W^*(G), E)$ be a D_G -valued random variable. Recall that the trivial D_G -valued n -th moments and cumulants of a are defined by

$$E(a^n) \quad \text{and} \quad k_n \left(\underbrace{a, \dots, a}_{n\text{-times}} \right).$$

In this chapter, we will deal with the following special D_G -valued random variable ;

Definition 7.1. Define an operator T in $W^*(G)$ by

$$(5.1) \quad T = \sum_{e \in E(G)} (L_e + L_e^*).$$

We will call T the generating operator of $W^*(G)$. The self-adjoint operators $L_e + L_e^*$, for $e \in E(G)$, are called the block operators of T .

Example 7.1. Let G be a one-vertex directed graph with N -edges. i.e.,

$$V(G) = \{v\} \quad \text{and} \quad E(G) = \{e_j = ve_jv : j = 1, \dots, N\}.$$

Then the graph W^* -algebra $W^*(G)$ satisfies that

$$(5.2) \quad W^*(G) = D_G *_{D_G} \left(\bigstar_{j=1}^N (W^*(\{L_{e_j}\}, D_G)) \right),$$

by Chapter 4. Notice that $D_G = \Delta_1 = \mathbb{C}$. Therefore, the formula (5.2) is rewritten by

$$(5.4) \quad W^*(G) = \bigstar_{j=1}^N (W^*(\{L_{e_j}\})),$$

where $*$ means the usual (scalar-valued) free product of Voiculescu. Also notice that $1_{D_G} = L_v = 1 \in \mathbb{C}$ and

$$(5.5) \quad L_{e_j}^* L_{e_j} = L_v = 1 = L_{e_j} L_{e_j}^*, \text{ for all } j = 1, \dots, N.$$

This shows that L_{e_j} 's are unitary in $W^*(G)$, for all $j = 1, \dots, N$. Now, define the generating operator $T = \sum_{j=1}^N (L_{e_j} + L_{e_j}^*)$ of $W^*(G)$. It is easy to see that each block operator $L_{e_j} + L_{e_j}^*$ is semicircular, by Voiculescu, for all $j = 1, \dots, N$. (Remember the construction of creation operators L_{e_j} 's and see [9].) Furthermore, by Chapter 3, we can get that all blocks $L_{e_j} + L_{e_j}^*$'s are free from each other in the graph W^* -probability space $(W^*(G), E)$.

By (5.4), the canonical conditional expectation $E : W^*(G) \rightarrow D_G$ is the faithful linear functional. Moreover, by (5.5), this linear functional E is a trace in the sense that $E(ab) = E(ba)$, for all $a, b \in W^*(G)$. From now, to emphasize that E is a trace, we will denote E by tr .

Let's compute the n -th cumulant of T ;

$$(5.6) \quad \begin{aligned} k_n(T, \dots, T) &= k_n \left(\sum_{j=1}^N (L_{e_j} + L_{e_j}^*), \dots, \sum_{j=1}^N (L_{e_j} + L_{e_j}^*) \right) \\ &= \sum_{j=1}^N k_n \left((L_{e_j} + L_{e_j}^*), \dots, (L_{e_j} + L_{e_j}^*) \right), \end{aligned}$$

by the mutual freeness of $\{L_{e_j}, L_{e_j}^*\}$'s on $(W^*(G), tr)$, for $j = 1, \dots, N$. Observe that

$$\begin{aligned}
& k_n \left((L_{e_j} + L_{e_j}^*), \dots, (L_{e_j} + L_{e_j}^*) \right) \\
(5.7) \quad & = \begin{cases} k_2 \left((L_{e_j} + L_{e_j}^*), (L_{e_j} + L_{e_j}^*) \right) & \text{if } n = 2 \\ 0 & \text{otherwise,} \end{cases}
\end{aligned}$$

by the semicircularity of $L_{e_j} + L_{e_j}^*$, for $j = 1, \dots, N$. By (5.7), the formula (5.6) is

$$\begin{aligned}
& k_n (T, \dots, T) \\
(5.8) \quad & = \begin{cases} \sum_{j=1}^N k_2 \left(L_{e_j} + L_{e_j}^*, L_{e_j} + L_{e_j}^* \right) & \text{if } n = 2 \\ 0 & \text{otherwise} \end{cases}
\end{aligned}$$

Now, observe $k_2 \left(L_{e_j} + L_{e_j}^*, L_{e_j} + L_{e_j}^* \right)$;

$$\begin{aligned}
& k_2 \left(L_{e_j} + L_{e_j}^*, L_{e_j} + L_{e_j}^* \right) \\
& = k_2 \left(L_{e_j}, L_{e_j} \right) + k_2 \left(L_{e_j}, L_{e_j}^* \right) + k_2 \left(L_{e_j}^*, L_{e_j} \right) + k_2 \left(L_{e_j}^*, L_{e_j}^* \right) \\
& = 0 + k_2 \left(L_{e_j}, L_{e_j}^* \right) + k_2 \left(L_{e_j}^*, L_{e_j} \right) + 0
\end{aligned}$$

by Section 2.1

$$= \text{tr} \left(L_{e_j} L_{e_j}^* \right) + \text{tr} \left(L_{e_j}^* L_{e_j} \right) = 2 \cdot \text{tr} \left(L_{e_j}^* L_{e_j} \right)$$

since tr is a trace

$$= 2 \cdot L_v = 2,$$

for $j = 1, \dots, N$, by Section 2.1 and 2.2. So, we can get that

$$(5.9) \quad k_n (T, \dots, T) = \begin{cases} 2N & \text{if } n = 2 \\ 0 & \text{otherwise.} \end{cases}$$

Now, we can compute the trivial moments of T , via the Möbius inversion.

$$\text{tr} (T^n) = \sum_{\pi \in NC(n)} k_\pi (a, \dots, a)$$

where $k_\pi(a, \dots, a) = \prod_{V \in \pi} k_{|V|} \left(\underbrace{a, \dots, a}_{|V|-\text{times}} \right)$, for each $\pi \in NC(n)$, by Nica and Speicher (See [1] and [17])

$$= \sum_{\pi \in NC_2(n)} k_\pi(a, \dots, a) = \sum_{\pi \in NC_2(n)} \left(\prod_{V \in \pi} k_{|V|}(a, a) \right)$$

where $NC_2(n) = \{\pi \in NC(n) : V \in \pi \Leftrightarrow |V| = 2\}$ is the collection of all noncrossing pairings

$$(5.10) \quad = \sum_{\pi \in NC_2(n)} \left(\prod_{V \in \pi} 2N \right) = \sum_{\pi \in NC_2(n)} (2N)^{|\pi|},$$

where $|\pi| \stackrel{\text{def}}{=} \text{the number of blocks in } \pi$. Notice that the above formula (5.10) shows us that the n should be even, because $NC_2(n)$ is nonempty when n is even. Therefore,

$$(5.11) \quad \text{tr}(T^n) = \begin{cases} \sum_{\pi \in NC_2(n)} (2N)^{|\pi|} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Also, notice that if $\pi \in NC_2(n)$, then $|\pi| = \frac{n}{2}$, for all even number $n \in \mathbb{N}$. So,

$$(5.12) \quad \begin{aligned} \text{tr}(T^n) &= \begin{cases} |NC_2(n)| \cdot (2N)^{\frac{n}{2}} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases} \\ &= \begin{cases} (2N)^{\frac{n}{2}} \cdot c_{\frac{n}{2}} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd,} \end{cases} \end{aligned}$$

where $c_k = \frac{1}{k+1} \binom{2k}{k}$ is the k -th Catalan number, for all $k \in \mathbb{N}$. Remember that

$$|NC(k)| = |NC_2(2k)| = c_k, \text{ for all } k \in \mathbb{N}.$$

Therefore, by (5.9) and (5.12), we can compute the moments and cumulants of the generating operator T of $(W^*(G), \text{tr})$;

$$\text{tr}(T^n) = \begin{cases} (2N)^{\frac{n}{2}} \cdot c_{\frac{n}{2}} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd,} \end{cases}$$

and

$$k_n(T, \dots, T) = \begin{cases} 2N & \text{if } n = 2 \\ 0 & \text{otherwise.} \end{cases}$$

Example 7.2. Let $N \in \mathbb{N}$ and let G be the circulant graph with

$$V(G) = \{v_1, \dots, v_N\}$$

and

$$E(G) = \{e_1, \dots, e_N\}$$

with

$$e_j = v_j e_j v_{j+1}, \text{ for } j = 1, \dots, N-1, \text{ and } e_N = v_N e_N v_1.$$

Define the generating operator $T = \sum_{j=1}^N (L_{e_j} + L_{e_j}^*)$ of the graph W^* -algebra $W^*(G)$. In this case, we can get the diagonal subalgebra D_G of $W^*(G)$, as a von Neumann algebra which is isomorphic to Δ_N , where Δ_N is a subalgebra of the matricial algebra $M_N(\mathbb{C})$. Define the canonical conditional expectation $E : W^*(G) \rightarrow D_G$. Then we can compute the trivial n -th D_G -valued cumulant of the operator T , by regarding it as a D_G -valued random variable in the graph W^* -probability space $(W^*(G), E)$ over $D_G = \Delta_N$. Notice that each block $L_{e_j} + L_{e_j}^*$'s are free from each other over D_G in $(W^*(G), E)$, by the diagram-distinctness of e_j 's, for $j = 1, \dots, N$.

Fix $n \in \mathbb{N}$. Then

$$\begin{aligned} k_n \left(\underbrace{T, \dots, T}_{n\text{-times}} \right) &= k_n \left(\sum_{j=1}^N (L_{e_j} + L_{e_j}^*), \dots, \sum_{j=1}^N (L_{e_j} + L_{e_j}^*) \right) \\ &= \sum_{j=1}^N k_n \left((L_{e_j} + L_{e_j}^*), \dots, (L_{e_j} + L_{e_j}^*) \right) \end{aligned}$$

by the mutual D_G -freeness of $\{L_{e_j}, L_{e_j}^*\}$'s, for $j = 1, \dots, N$

$$(5.13) \quad = \sum_{j=1}^N \sum_{(u_1, \dots, u_n) \in \{1, *\}^n} k_n \left(L_{e_j}^{u_1}, \dots, L_{e_j}^{u_n} \right).$$

Recall that, by Section 2.2, we can get that

$$(5.14) \quad k_n \left(L_{e_j}^{u_1}, \dots, L_{e_j}^{u_n} \right) = \mu_{e_j, \dots, e_j}^{u_1, \dots, u_n} \cdot \text{Pr oj} \left(L_{e_j}^{u_1} \dots L_{e_j}^{u_n} \right),$$

$$\text{where } \mu_{e_j, \dots, e_j}^{u_1, \dots, u_n} = \sum_{\pi \in C_{e_j, \dots, e_j}^{u_1, \dots, u_n}} \mu(\pi, 1_n).$$

Observe that since e_j 's are non-loop edges, $e_j^k \notin \mathbb{F}^+(G)$, for all $k \in \mathbb{N} \setminus \{1\}$, for $j = 1, \dots, N$. In other words, such e_j^k is not admissible. So, if $(u_1, \dots, u_n)$ is not alternating, in the sense that $(u_1, \dots, u_n) = (1, *, \dots, 1, *)$ or $(*, 1, \dots, *, 1)$, then $\text{Proj} \left(L_{e_j}^{u_1} \dots L_{e_j}^{u_n} \right) = 0_{D_G}$. For instance, $E \left(L_{e_j}^* L_{e_j} L_{e_j}^* \right) = 0_{D_G}$ or $E \left(L_{e_j}^2 L_{e_j}^* L_{e_j} \right) = 0_{D_G}$, by Section 2.1. Therefore, the only nonvanishing case is either

$$k_n \left(L_{e_j}, L_{e_j}^*, \dots, L_{e_j}, L_{e_j}^* \right) \quad \text{or} \quad k_n \left(L_{e_j}^*, L_{e_j}, \dots, L_{e_j}^*, L_{e_j} \right),$$

where n is even. Notice that

$$(5.15) \quad \mu_{e_j, e_j, \dots, e_j, e_j}^{1, *, \dots, 1, *} = \mu_{e_j, e_j, \dots, e_j, e_j}^{*, 1, \dots, *, 1},$$

because $C_{e_j, e_j, \dots, e_j, e_j}^{1, *, \dots, 1, *} = C_{e_j, e_j, \dots, e_j, e_j}^{*, 1, \dots, *, 1}$, for all $j = 1, \dots, N$. Moreover, since $C_{e_j, e_j, \dots, e_j, e_j}^{1, *, \dots, 1, *} = C_{e_k, e_k, \dots, e_k, e_k}^{1, *, \dots, 1, *}$, for all $j \neq k$ in $\{1, \dots, N\}$,

$$(5.16) \quad \mu_{e_j, e_j, \dots, e_j, e_j}^{1, *, \dots, 1, *} = \mu_{e_k, e_k, \dots, e_k, e_k}^{1, *, \dots, 1, *},$$

for all $j, k \in \{1, \dots, N\}$. Let's denote $\mu_{e_j, e_j, \dots, e_j, e_j}^{1, *, \dots, 1, *}$ by μ_n , for all $j = 1, \dots, N$. Then, by (5.14), we have that

$$(5.17) \quad k_n \left(L_{e_j}^{u_1}, \dots, L_{e_j}^{u_n} \right) = \begin{cases} \mu_n L_{v_j} & \text{if } (u_1, \dots, u_n) = (1, *, \dots, 1, *) \\ \mu_n L_{v_{j+1}} & \text{if } (u_1, \dots, u_n) = (*, 1, \dots, *, 1) \\ 0_{D_G} & \text{otherwise,} \end{cases}$$

for all $j = 1, \dots, N$, where $L_{v_{N+1}}$ means L_{v_1} . So, by (5.13) and (5.17), we can get that

$$\begin{aligned} & k_n(T, \dots, T) \\ &= \sum_{j=1}^N \left(k_n \left(L_{e_j}, L_{e_j}^*, \dots, L_{e_j}, L_{e_j}^* \right) + k_n \left(L_{e_j}^*, L_{e_j}, \dots, L_{e_j}^*, L_{e_j} \right) \right) \\ &= \sum_{j=1}^N (\mu_n L_{v_j} + \mu_n L_{v_{j+1}}) = \sum_{j=1}^N \mu_n (L_{v_j} + L_{v_{j+1}}) \end{aligned}$$

where $L_{v_{N+1}}$ means L_{v_1} , for all $n \in 2\mathbb{N}$. Therefore,

$$k_n(T, \dots, T) = \begin{cases} \sum_{j=1}^N \mu_n (L_{v_j} + L_{v_{j+1}}) & \text{if } n \text{ is even} \\ 0_{D_G} & \text{if } n \text{ is odd.} \end{cases}$$

$$(5.18) \quad = \begin{cases} 2\mu_n \cdot 1_{D_G} & \text{if } n \text{ is even} \\ 0_{D_G} & \text{if } n \text{ is odd.} \end{cases}$$

Unfortunately, it is very hard to compute μ_n , when $n \rightarrow \infty$. But we have to remark that if we have arbitrary graph H and its graph W^* -probability space $(W^*(H), F)$ over its diagonal subalgebra D_H and if $w \in \text{loop}^c(G)$, then

$$\mu_{w, w, \dots, w, w}^{1, *, \dots, 1, *} = \mu_n = \mu_{w, w, \dots, w, w}^{*, 1, \dots, *, 1}, \text{ for all } n \in 2\mathbb{N}.$$

Now, let's compute the trivial n -th D_G -valued moment of T . Notice that since all odd trivial D_G -valued cumulants of T vanish, all odd trivial D_G -valued moments of T vanish (See [11] and [14]). Thus it suffices to compute the even trivial D_G -valued moments of T . Assume that $n \in 2\mathbb{N}$. Then

$$(5.19) \quad E(T^n) = \sum_{\pi \in NC_E(n)} k_\pi(T, \dots, T),$$

where $k_\pi(T, \dots, T)$ is the partition-dependent cumulant of T (See [16]) and

$$NC_E(n) \stackrel{\text{def}}{=} \{\pi \in NC(n) : V \in \pi \Leftrightarrow |V| \in 2\mathbb{N}\}.$$

By (5.18), we can get that $k_n(T, \dots, T)$ commutes with all elements in $W^*(G)$, because 1_{D_G} and 0_{D_G} commutes with $W^*(G)$ and $2\mu_n \in \mathbb{C}$, for all $n \in \mathbb{N}$. So, the formula (5.19) can be reformed by

$$\begin{aligned} E(T^n) &= \sum_{\pi \in NC_E(n)} \left(\prod_{V \in \pi} k_{|V|}(T, \dots, T) \right) \\ &= \sum_{\pi \in NC_E(n)} \left(\prod_{V \in \pi} 2\mu_{|V|} \cdot 1_{D_G} \right) \\ (5.20) \quad &= \left(\sum_{\pi \in NC_E(n)} \left(\prod_{V \in \pi} 2\mu_{|V|} \right) \right) \cdot 1_{D_G}, \end{aligned}$$

for all $n \in 2\mathbb{N}$. Therefore, by (5.18) and (5.20), we have that if T is the generating operator of the graph W^* -algebra of the circulant graph G with N -vertices, then

$$E(T^n) = \begin{cases} \left(\sum_{\pi \in NC_E(n)} \left(\prod_{V \in \pi} 2\mu_{|V|} \right) \right) \cdot 1_{D_G} & \text{if } n \text{ is even} \\ 0_{D_G} & \text{if } n \text{ is odd.} \end{cases}$$

and

$$k_n \left(\underbrace{T, \dots, T}_{n\text{-times}} \right) = \begin{cases} 2\mu_n \cdot 1_{D_G} & \text{if } n \text{ is even} \\ 0_{D_G} & \text{if } n \text{ is odd.} \end{cases}$$

Reference

- [1] A. Nica, R-transform in Free Probability, IHP course note, available at www.math.uwaterloo.ca/~anica.
- [2] A. Nica and R. Speicher, R-diagonal Pair-A Common Approach to Haar Unitaries and Circular Elements, (1995), www.mast.queensu.ca/~speicher.
- [3] B. Solel, You can see the arrows in a Quiver Operator Algebras, (2000), preprint
- [4] A. Nica, D. Shlyakhtenko and R. Speicher, R-cyclic Families of Matrices in Free Probability, J. of Funct Anal, 188 (2002), 227-271.
- [5] D. Shlyakhtenko, Some Applications of Freeness with Amalgamation, J. Reine Angew. Math, 500 (1998), 191-212.
- [6] D. Voiculescu, K. Dykemma and A. Nica, Free Random Variables, CRM Monograph Series Vol 1 (1992).
- [7] D. Voiculescu, Operations on Certain Non-commuting Operator-Valued Random Variables, Astérisque, 232 (1995), 243-275.
- [10] D. Shlyakhtenko, A-Valued Semicircular Systems, J. of Funct Anal, 166 (1999), 1-47.
- [10] D.W. Kribs and M.T. Jury, Ideal Structure in Free Semigroupoid Algebras from Directed Graphs, preprint
- [10] D.W. Kribs and S.C. Power, Free Semigroupoid Algebras, preprint
- [11] I. Cho, Amalgamated Boxed Convolution and Amalgamated R-transform Theory, (2002), preprint.
- [12] I. Cho, The Tower of Amalgamated Noncommutative Probability Spaces, (2002), Preprint.
- [13] I. Cho, Free Perturbed R-transform Theory, (2003), Preprint.
- [14] I. Cho, Compatibility of a Noncommutative Probability Space and a Noncommutative Probability Space with Amalgamation, (2003), Preprint.
- [15] I. Cho, Graph W^* -Probability Spaces Over the Diagonal Subalgebras, (2004), Preprint.
- [16] R. Speicher, Combinatorial Theory of the Free Product with Amalgamation and Operator-Valued Free Probability Theory, AMS Mem, Vol 132 , Num 627 , (1998).
- [17] R. Speicher, Combinatorics of Free Probability Theory IHP course note, available at www.mast.queensu.ca/~speicher.
- [18] T. Bates and D. Pask, Flow Equivalence of Graph Algebras, (2004), Preprint.